

ORIGINAL RESEARCH ARTICLE

Assessment of Serological Markers of Genital *Chlamydia trachomatis* Infection Among the Gynaecology Patients attending Babcock University Teaching Hospital, Ilishan-Remo, Ogun State, Nigeria

DOI: 10.29063/ajrh2019/v23i4.7

Tinuade A Ajani^{1*}, Charles J. Elikwu¹, Victor Nwadike¹, Tayo Babatunde¹, Chinenye G Anaedobe², Opeoluwa Shonekan¹, Chika Okangba¹, Timothy Oluwasola³, Azubuike Omeonu¹, Bibitayo Faluyi¹, Tunde E Thompson¹, Ejime Ebeigbe¹, Mustapha A Ajani⁴, Amelia K Joshua¹, Titilope Kolawole¹, Heritage Kristilere¹, Chibuikwe M Meremikwu¹, Lucky Mgbemena¹, Chika S Nwaejike¹, Ayodeji Salami¹, Anatorun Tantua¹, Mayowa Timothy¹, Tobenna Akagbusum¹ and Akintoye Ol Coker¹

Department of Medical Microbiology, Babcock University Teaching Hospital, Ilishan-Remo, Ogun State, Nigeria¹; Department of Medical Microbiology, University of Abuja, Federal Capital territory, Abuja²; Department of Obstetrics and Gynaecology, University College Hospital, Ibadan, Oyo State, Nigeria³; Department of Histopathology, Babcock University Teaching Hospital, Ilishan-Remo, Ogun State, Nigeria⁴

*For Correspondence: Email: solamustoo@yahoo.com; Phone: +2348034412609

Abstract

Genital *Chlamydia trachomatis* infection causes significant morbidity and mortality in women. A number of epidemiologic studies have suggested that Polymerase Chain Reaction (PCR) is more accurate as a diagnostic tool for *Chlamydia trachomatis*. However, the use of serological markers may be cost effective and practical in diagnosing and estimating the burden of the disease in resource limited countries. This study was aimed at determining the serological markers (IgG, IgM and IgA) of *Chlamydia trachomatis*, evaluate the association between *Chlamydia trachomatis* infection and the sociodemographic characteristics and clinical diagnosis of the participants. This was a cross sectional hospital-based study in which blood samples from 145 consenting participants were tested for IgG, IgM and IgA antibodies against *Chlamydia trachomatis* using enzyme linked immunosorbent assay and their clinical diagnosis, retrieved from their case notes. The cumulative prevalence of seropositivity for *Chlamydia trachomatis* (IgG, IgM, IgA) was 112 (77.2%) while 33 (22.8%) were seronegative. The overall predominant seromarker was IgG 91(62.8%) while IgM and IgA accounted for 85(58.6%) and 54(37.2%) respectively. A statistically significant association was found between *Chlamydia trachomatis* infection and PID (p value = 0.031), primary infertility (p value 0.011) and level of income (p value= (0,045). (*Afr J Reprod Health* 2019; 23[4]: 54-62).

Keywords: *Chlamydia trachomatis*, Serological markers IgG, IgM, IgA, PID

Résumé

L'infection génitale à *Chlamydia trachomatis* provoque une morbidité et une mortalité importantes chez les femmes. Un certain nombre d'études épidémiologiques ont suggéré que la réaction en chaîne par polymérase (RCP) est plus précise en tant qu'outil de diagnostic pour *Chlamydia trachomatis*. Cependant, l'utilisation des marqueurs sérologiques peut être rentable et pratique pour diagnostiquer et estimer la charge de la maladie dans les pays à ressources limitées. Cette étude visait à déterminer les marqueurs sérologiques (IgG, IgM et IgA) de *Chlamydia trachomatis*, évaluer l'association entre l'infection à *Chlamydia trachomatis* et les caractéristiques sociodémographiques et le diagnostic clinique des participantes. Il s'agissait d'une étude transversale en milieu hospitalier dans laquelle des échantillons de sang de 145 participantes consentantes ont été testés pour les anticorps IgG, IgM et IgA contre *Chlamydia trachomatis* en utilisant un dosage immuno-enzymatique et leur diagnostic clinique, extraits de leurs notes de cas. La prévalence cumulative de la séropositivité pour *Chlamydia trachomatis* (IgG, IgM, IgA) était de 112 (77,2%) tandis que 33 (22,8%) étaient séronégatives. Le séromarqueur prédominant global était l'Ig G 91 (62,8%) tandis que l'Ig M et l'IgA représentaient respectivement 85 (58,6%) et 54 (37,2%). Une association statistiquement significative a été trouvée entre l'infection à *Chlamydia trachomatis* et le PID (valeur p = 0,031), l'infertilité primaire (valeur p 0,011) et le niveau de revenu (valeur p = (0,045). (*Afr J Reprod Health* 2019; 23[4]: 54-62).

Mots-clés: *Chlamydia trachomatis*, marqueurs sérologiques IgG, IgM, IgA, PID

Introduction

Chlamydia trachomatis serovars D through K typically cause cervicitis, PID in women and non-gonococcal urethritis in men¹. Prevalence is consistently highest among young women aged 15 to 24 years compared to others and higher among blacks than other races². Chlamydial infection in females is often asymptomatic or present with less severe symptoms than other sexually transmitted diseases. The asymptomatic nature precludes early diagnosis and treatment. This ultimately leads to complications such as Pelvic inflammatory diseases (PID) as chronic pelvic pain sets in, ectopic pregnancy, salpingitis, recurrent abortions and infertility^{1,3-4}.

According to the World Health Organization (WHO), About 131 million people are infected with Chlamydia each year⁵. Centre for Disease Control and prevention estimated approximately 3 to 4 million new cases of Chlamydia infection in the United States each year and 75% of such new cases are diagnosed among asymptomatic women⁶.

In Nigeria, the prevalence of Chlamydia cervicitis was found to be 17.6% among women with infertility³ while another study reported a prevalence of 33% among asymptomatic women⁷. A study also reported a prevalence of 7.3% among asymptomatic infertile women in Ibadan⁸.

Considering the burden of Chlamydia, the United States Centre for Disease Control and Prevention recommends annual screening of women of reproductive age to ensure early diagnosis and treatment. This has been implemented in developed countries, but is yet to be done in developing countries like Nigeria^{9, 10}. Unless implemented, the infection will remain a "silent epidemic"¹¹.

Although advances in highly accurate and non-invasive diagnostic testing such as PCR have allowed for better estimation of the burden of the disease, particularly the asymptomatic state, however, these methods of diagnosis need specialized equipment, infrastructure and trained personnel. Thus they are neither cost effective nor readily accessible in resource limited countries like Nigeria¹².

Considering the high prevalence of 33% among asymptomatic women as reported in a study in Nigeria⁷, serological diagnosis may be more practical and accessible when compared to the molecular assays for *Chlamydia trachomatis*. While most patients are likely to present when there is a complication, a positive serology result on routine screening of at risk patients may be the only indication of chlamydial involvement¹³.

The higher sensitivity and specificity of qualitative third generation ELISA for *Chlamydia trachomatis* serological markers makes it more reliable than the rapid diagnostic test strips used in most laboratories. Our facility uses the rapid tests for chlamydia diagnosis; this often yields negative results which are inaccurate and misleading.

Therefore, the aim of this study is to determine the serological markers of genital *Chlamydia trachomatis* among patients presenting at the gynaecology clinic of the Babcock University teaching Hospital, Ilishan-Remo, Ogun State, Nigeria and to evaluate the association between *Chlamydia trachomatis* infection and the sociodemographic characteristics and clinical diagnosis of the participants.

Methods

Study design

This was a cross sectional hospital-based study conducted in the gynaecology clinic of Babcock University Teaching Hospital from November 2017 to July 2018.

Study population

The participants were gynaecology patients attending the clinic at this period.

Inclusion criteria

Consenting females within the reproductive age, who were willing to fill the questionnaire, allow blood collection for *Chlamydia trachomatis* testing were included in the study.

Exclusion criteria

Those excluded were females who were on antibiotics or who had used antibiotics within the

previous 6 months and those who were not willing to provide consent.

Sample size determination

A sample size of 145 was calculated based on *Chlamydia trachomatis* seroprevalence of 9.2% found among gynaecology and STI patients in Kano, Northern part of Nigeria, to give a 95% confidence level and margin of error of $\pm 5\%$.

Specimen collection and processing

Blood samples were tested using qualitative third generation enzyme linked immunosorbent assay (ELISA), type-specific for IgG, IgM and IgA against polypeptide derived from *Chlamydia trachomatis* major outer-membrane antigen (MOMP) (DIAPRO Diagnostic Bioprobes Milano Italy). All the blood samples were tested separately for each of these serology markers.

Data collection procedure

Pretested interviewer-based structured questionnaires were used to obtain the sociodemographic factors of the women and clinical diagnosis retrieved from their case notes.

Data analysis

Standard descriptive and inferential statistical analysis was carried out using SPSS version 23. (SPSS Inc. Illinois, USA). Means and standard deviations were derived for quantitative variables while proportions were derived for qualitative variables. Association between categorical variables was determined using Chi-square test at statistical significance level of set 5%. Ethical approval was received from Babcock University ethical review committee.

Results

The age of the 145 participants ranged from 15 to 45 years with the mean age of 23.7 years (SD 1.483). Majority of the participants were within the age range of 15 to 20 years 70(48.3%). Majority 127 (87.6%) of them were students, 110 (75.9%) were unmarried and 119 (82.1%) had

tertiary education. The socio-demographic characteristics are illustrated in Table 1.

Majority 44 (30.3%) of the participants had PID, 22 (15.2%) had Dysmenorrhea, 14 (9.7%) had primary infertility, 11 (7.6%) was diagnosed with cervicitis, 7 (4.8%) had Secondary infertility, 6 (4.1%) had recurrent abortion and 41 (28.3%) were asymptomatic women who presented for preschool medical checkup.

Of the 145 participants, the cumulative seroprevalence for *Chlamydia trachomatis* (IgG, IgM, IgA) was 112(77.2%) while 33 (22.8%) were seronegative. Of the 112 seropositive participants, 37 (33%) had one seromarker, 32(28.6%) had two seromarkers, while 43 (38.4%) were positive for all three markers. Of the 37 who had one seromarker, 21(56.8%) were IgG, 13(35.1%) were IgM and 3(8.1%) were IgA. Of the 32 participants positive for 2 seromarkers, 24(75%) had both IgG and IgM, 5(15.6%) had IgM and IgA, while 3(9.4%) had both IgG and IgA.

In total, 91 (62.8%) participants were positive for IgG making it the predominant seromarker, 85(58.6%) were positive for IgM and 54 (37.2%) for IgA. The overall combined prevalence of *Chlamydia trachomatis* IgG and IgA was 46(31.7%), IgM and IgA was 48 (33.1%), IgG and IgM was 67 (46.2%). This is further illustrated in Table 2.

The relationship between *Chlamydia trachomatis* serological markers and the sociodemographic characteristics as well as clinical diagnosis of the participants are shown in tables 3 and 4 respectively. Table 5 shows the association between *Chlamydia trachomatis* infection and the sociodemographic and clinical diagnosis of the participants.

Discussion

Molecular diagnostic tools for *Chlamydia trachomatis* infection such as PCR has allowed for more accurate diagnosis and estimation of the disease burden but in resource limited countries such as ours, molecular diagnosis is expensive and not routinely found in most laboratories. Readily available screening tools such as serological markers are more practicable for determination of *Chlamydia trachomatis* infection.

Table1: Sociodemographic characteristics of the participants (N=145)

Variables	Frequency	Percentage
Age		
15-20	70	48.3
21-25	35	24.1
26-30	15	10.3
31-35	8	5.5
36-40	10	6.9
41-45	7	4.8
Marital status		
Single	110	75.9
Married	32	22.1
Widowed	2	1.4
Separated	1	0.7
Level of Education		
No schooling	3	2.1
Primary	3	2.1
Secondary	20	13.8
Tertiary	119	82.1
Type of housing		
Personal	88	60.7
Rented(\geq 2bedrooms)	29	20.0
Self-contained	16	11.0
Single room	7	4.8
Others	5	3.4
Level of income (Naira)		
< 18000	75	51.8
\geq 18000	70	48.2
Occupation		
Students	127	87.6
Professional	4	2.8
Artisan	4	2.8
Non-professional	10	6.9

In the index study, the cumulative prevalence of *chlamydia trachomatis* serologic markers was 77.2%. This is similar to the cumulative prevalence of 68.5% reported by Joyee AG et al in Indian where the three serology markers of *chlamydia trachomatis* were utilized as well as prevalence of 70.8% reported by Moses *et al* in Ilorin and 74% among subfertile women in Port Harcourt both in Nigeria¹³⁻¹⁴. While it is higher than rates reported in other regions in Nigeria such as 56.1% among gynaecology patients in Jos by Mawak et al¹⁶ and 51% among pregnant women in Lagos state by Okoror et al¹⁷, it is at variance with other studies in that reported much lower prevalence rates¹⁸⁻¹⁹. These variances in the prevalence rates across the country can be attributed to varying study population, sensitivity and specificity of laboratory diagnostic tool used.

The highest prevalence of *Chlamydia trachomatis* was observed among women in the 15- 25 age group 105 (72.4%), students 127 (87.6%), and those with tertiary level education 119 (82.1%). This is in line with previous studies in Nigeria and Argentina where a high prevalence rate was reported among young people^{3, 20, 21}. These findings may suggest high sexual activity, lack or inconsistent use of condom, inadequate sexual health education and multiple sex partners among young women²². Furthermore, chlamydia infection was significantly higher amongst the unmarried women compared to the married ones. This is in agreement with previous finding reported in Enugu¹⁸ but in contrast to data reported in Jos¹⁶ both in Nigeria. This comparison however, should be done with caution as majority of the participants in our study were unmarried.

Majority of our study participants presented with PID 44(30.3%) and 91% were positive for at least one of the sero markers (IgG, IgM, IgA). Our finding is consistent with other studies carried out in India and America^{13,23}. This observation suggests that serological studies may be useful in identifying *Chlamydia trachomatis* in ascending upper genital tract infection. Moreover, some studies have demonstrated that serology has been helpful in cases where direct test such as PCR, has failed to detect the presence of *Chlamydia trachomatis* in PID patients^{13,23-24}. Indeed, serology prevents the morbidity associated with invasive procedures such as laparoscopy and hysterosalpingography (HSG) in women presenting with *Chlamydia trachomatis* sequelae²⁴.

Type specific distribution shows that IgG seromarker was the most predominant (62.8%) and its presence indicates chronic infection or previous exposure. Seromarker IgM (58.6%), the second predominant marker in our study, signifies the presence of acute infection. However a previous study observed that antibodies produced during *chlamydia trachomatis* acute genital infection are usually long lived and cannot distinguish between current and previous infection²⁵. Seromarker IgA (37.2%), the least predominant in our study has been reported as a better indicator of active infection because it has a half live of 5-7 days²⁶⁻²⁷, thus useful for monitoring active infection after

Table 2: Cumulative prevalence of *chlamydia trachomatis* serological markers among the participants (N=145)

<i>Chlamydia trachomatis</i> IgG	<i>Chlamydia trachomatis</i> IgM		
	Positive (%)	Negative (%)	Total (%)
Positive	67 (46.2%)	24 (16.6%)	91 (62.8%)
Negative	18 (12.4%)	36 (24.8%)	54 (37.2%)
Total	85 (58.6%)	60 (41.4%)	145 (100%)
<i>Chlamydia trachomatis</i> IgG	<i>Chlamydia trachomatis</i> IgA		
	Positive (%)	Negative (%)	Total (%)
Positive	46 (31.7%)	45 (31.1%)	91 (62.8%)
Negative	8 (5.5%)	46 (31.7%)	54 (37.2%)
Total	54 (37.2%)	91 (62.8%)	145 (100%)
<i>Chlamydia trachomatis</i> IgM	<i>Chlamydia trachomatis</i> IgA		
	Positive (%)	Negative (%)	Total (%)
Positive	48 (33.1%)	37 (25.6%)	85 (58.6%)
Negative	6 (4.1%)	54 (37.2%)	60 (41.4%)
Total	54 (7.2%)	91 (62.8%)	145 (100%)

Table 3: Relationship between *Chlamydia trachomatis* serological markers and Sociodemographic factors of the Participants

Variables	<i>Chlamydia trachomatis</i> IgG		<i>Chlamydia trachomatis</i> IgM		<i>Chlamydia trachomatis</i> IgA	
	Positive	%	Positive	%	Positive	%
Age						
15-20	40	43.9	33	38.8	22	40.7
21-25	27	29.7	24	28.2	12	22.2
26-30	8	8.8	10	11.8	8	14.8
30-35	4	4.4	6	7.1	4	7.4
36-40	8	8.8	8	9.4	5	9.3
40-45	4	4.4	4	4.7	3	5.6
Total	91	100	85	100.0	54	100.0
Marital status						
Single	70	76.9	61	71.8	37	68.5
Married	19	20.9	22	25.9	16	29.6
Widowed	1	1.1	1	1.2	1	1.9
Separated	1	1.1	1	1.2	0	0.0
Total	91	100.0	85	100.1	54	100
Occupation						
Student	80	87.9	77	90.6	48	88.9
Professional	2	2.2	2	2.4	1	1.9
Artisan	2	2.2	2	2.4	1	1.9
Non- Professional	7	7.7	4	4.7	4	7.4
Total	91	100	85	100	54	100
Level of Education						
No schooling	2	2.3	2	2.4	2	3.7
Primary	3	3.3	1	1.2	0	0
Secondary	15	16.5	15	17.7	10	18.5
Tertiary	71	78.0	67	78.8	42	77.8
Total	91	100	85	100	54	100
Type of housing						
Personal	54	59.3	48	56.5	29	53.7
Rented	19	20.9	19	22.4	14	25.9
Self- contained	11	12.1	10	11.8	7	12.9
Single room	6	6.6	5	5.9	3	5.6
Others	1	1.1	3	3.5	1	1.9
Total	91	100	85	100	54	100




Figure 1: Clinical diagnosis of the participants N=145

Table 4: Relationship between Chlamydia trachomatis serological markers and clinical diagnosis of the participants (N=145)

Variables	Chlamydia trachomatis IgG		Chlamydia trachomatis IgM		Chlamydia trachomatis IgA	
	Positive	%	Positive	%	Positive	%
Clinical Diagnosis						
PID	32	35.2	33	38.8	14	25.9
Primary infertility	7	7.7	5	5.9	6	11.1
Secondary infertility	3	3.3	3	3.5	3	5.6
Recurrent abortion	4	4.4	3	3.5	2	3.7
Cervicitis	7	7.7	7	8.2	5	9.3
Dysmenorrhea	12	13.2	10	11.8	9	16.7
Assymptomatic	26	28.6	24	28.3	15	27.8
Total	91	100.	85	100	54	100.

treatment²⁶ In resource limited countries where access to molecular tests such as PCR is limited, seromarkers may be the only indicator that provides evidence of patients contact with *Chlamydia trachomatis*.

Some studies has suggested that the presence of IgG and IgA is a better indicator of chronic or persistent infection^{25,28}. Our study shows the overall prevalence of chronic/ persistent infection was 46 (31.7%) based on the presence of *Chlamydia trachomatis* IgG and IgA altogether.

This finding suggest that a high percentage of these women have chronic persistent chlamydia infection which can be transferred to their sexual partners or if /when pregnant, to their newborn thereby worsening the silent epidemic.

Cofactors significantly associated with increased positivity of *Chlamydia trachomatis* infection were PID, primary infertility and low level of income. This further buttresses the need for use of cost effective diagnostic tool to routinely screen women of reproductive age

Table 5: Association of chlamydia trachomatis infection with Socio-demographic factors and clinical diagnosis

Variables	Chlamydia infection		X ²	P -value
	Positive(%)	Negative		
Sociodemographic factors				
Age (years)				
15-20	50(71.4)	20(28.6)	5.339	0.378
21-25	28(80.0)	7 (20.0)		
26-30	11(73.3)	4 (26.7)		
31-35	7 (87.5)	1 (12.5)		
36-40	10(100.0)	0 (0.0)		
41-45	6 (85.7)	1 (14.3)		
Marital status				
Single	83(75.5)	27(24.5)	2.265	0.159
Married	27(84.4)	5 (15.6)		
Widowed	1 (50.0)	1 (50.0)		
Separated	1 (100.0)	0 (0.0)		
Level of Education				
No schooling	2 (66.7)	1 (33.3)	1.202	0.753
Primary	3 (100.0)	0 (0.0)		
Secondary	16 (80.0)	4 (20.0)		
Tertiary	91 (76.5)	28 (23.5)		
Type of housing				
Personal	65 (73.9)	23(26.1)	2.250	0.690
Rented (≥ 2)	25 (86.2)	4 (13.8)		
Self-contained	12(75.0)	4 (25.0)		
Single room	6(85.7)	1(14.3)		
Others	4(80.0)	1 (20.0)		
Level of income (Naira)				
< 18000	63(84.0)	12(16.0)	4.037	0.045
≥18000	49(70.0)	21(30.0)		
Occupation				
Students	99(78.0)	28(22.0)	1.780	0.619
Professionals	2 (50.0)	2 (50.0)		
Artisans	3 (75.0)	1(25.0)		
Non- professionals	8 (80.0)	2(20.0)		
Clinical diagnosis				
PID	39 (88.6)	5(11.4)	4.666	0.031
Primary infertility	7 (50.0)	7(50.0)	6.439	0.011
Secondary infertility	3(42.9)	4(57.1)	4.947	0.26
Recurrent abortion	4(66.7)	2(33.3)	0.398	0.528
Cervicitis	9(81.8)	2(18.2)	0.142	0.706
Dysmenorrhea	17(77.3)	5(22.7)	0.000	0.997
Asymptomatic	33(80.5)	8 (19.5)	0.343	0.558
P value less than 0.05 is taken as significant				

particularly those at risk and thus aid early diagnosis, treatment and prevent morbidity associated with *Chlamydia trachomatis* infection. This will go a long way in reducing the burden of the disease and address the silent epidemic.

Conclusion

Use of serologic markers is practical and cost effective in diagnosing *Chlamydia trachomatis* infection in symptomatic and asymptomatic

women and is a useful tool for routine use in screening women at risk for the infection. In view of the potential sequelae and the morbidity associated with *Chlamydia trachomatis* among females, it may be necessary to implement prevention and control strategies like screening programmes by using serological markers. This is even more pertinent in resource limited country like ours where molecular diagnostic tools like PCR might not be affordable, accessible and therefore impracticable.

Contribution of Authors

TINUADE ADESOLA AJANI contributed to this article by conception, design, acquisition, analysis and interpretation of data, drafting the article, final approval of the version to be published and general supervision of the work. CHARLES JOHN ELIKWU, VICTOR UGOCHUKWU NWADIKE, BABATUNDE TAYO, OPEOLUWA SHONEKAN, CHIKA CELEN OKANGBA, CHINENYE GLORIA ANAEDOBE, TUNDE EHIMEN THOMPSON, EJIME EBEIGBE, AZUBUIKE OMEONU, BIBITAYO FALUYI contributed to this article by the design and interpretation of the data, drafting the article and final approval of the version to be published. MUSTAPHA AKANJI AJANI, TIMOTHY OLUWASOLA AKINTOYE OLUSEGUN COKER contributed by analysis and interpretation of data, drafting the article and final approval of the version to be published while AMELIA KIKI- JOSHUA, TITILOPE KOLAWOLE, HERITAGE KRISTILERE, CHIBUIKE MARTIN MEREMIKWU, LUCKY MGBEMENA, CHIKA SOLOMON-NWAEJIKE, AYODEJI SALAMI, ANATORUN TANTUA, MAYOWA TIMOTHY, TOBENNA AKAGBUSUM contributed to the acquisition, analysis and interpretation of data and drafting the article.

References

- Siemer J, Theile O, Larbi Y, Fasching PA, Danso KA, Kreienberg R and Essig A. Chlamydia trachomatis infection as a risk factor for infertility among women in Ghana, West Africa. *Am. J. Trop. Med. Hyg.* 2008; 78(2):323–327.
- Otoikhian CS, Okoror LE and Ekakitie AO. Polymerase Chain Reaction: a genetic tool for assessing Chlamydia trachomatis in mid-stream urine of asymptomatic female. *Int J life Scien. Biotechnol. Pharm research* 2012; 1(4):213-221.
- Atalabi OM, Fayemiwo SA, Oladokun AA and Bakare RA. Pattern of Asymptomatic Sexually transmitted infections in Women undergoing hysterosalpingography for infertility evaluation in Ibadan Nigeria. *Trop. J. Obstet Gynaecol* 2013; 30(2):91-98.
- Nsonwu-Anyanwu AC, Charles-Davies MA, Oni AA, Taiwo VO and Bello FA. chlamydial infection, plasma peroxidation and obesity in tubal Infertility. *Ann. Ibd.pg Med* 2011; 9(2):82-88
- World Health Organisation: WHO Library Cataloguing in Publication Data: report on global sexually transmitted infection surveillance 2015. World Health Organisation Switzerland, 2016S:3-5
- Molicotti P, Usai D, Cubeddu M, Sechi LA and Zanetti S. Comparison of two molecular methods for diagnosis of Chlamydia trachomatis. *J Infect Dev Ctries.* 2013 Jan 15; 7(1):64-66
- Isibor JO, Ugbomoiko D, Nwobu GO, Ekundayo AO, Eweani IB and Okogun GRA. Detection of Chlamydial antigen in cervical specimens from antenatal clinic in Benin City, Nigeria. *Af J Clin Exp Microbiol.* 2005; 6:208-211.
- Ajani TA, Fayemiwo SA, Oluwasola TA, Anaedobe CG, Ajani MA and Bakare RA. Prevalence of Asymptomatic genital Chlamydia trachomatis infection among infertile women in Ibadan, Nigeria using Polymerase Chain Reaction. *Indian J. Med. Research and Pharmaceutical Sciences.* 2017; 4(9): 13-24
- Center for Disease Control and Prevention. Sexually transmitted disease surveillance 2018: Chlamydia [Cited 18th November, 2019]. Available from :<https://www.cdc.gov/std/stats18/chlamydia.html>
- Center for Disease Control and Prevention. Sexually transmitted Diseases: STD and HIV Screening recommendation. CDC A-Z index. [Cited on the 23rd October, 2018} Available from: <https://www.cdc.gov/std/prevention/screeningrecs.htm>
- Okoror LE, Agbonlahor DE and Umolu PI. Prevalence of chlamydia in patients attending gynaecology clinic in South Eastern Nigeria. *African Health Science.* 2007; 7(1): 18-24
- Rashidi BH, Chamani-Tabriz L, Haghollahi F, Jeddi-Tehrani M, Naghizadeh MM, Shariat M, Akhondi MM, Bagheri R, Asgari S, and Wylie K. Effects of Chlamydia trachomatis infection on fertility, a case control study. *J Reprod. Fertility* 2013; 14(55):67-72.
- Joyee AG, Thyagarajan SP, Vikram Reddy E, Rajendran P, Venkatesan C and Ganapathy M. Diagnostic utility of serologic markers for genital chlamydial infection in STD patients in Chennai India. *J Assoc Physicians India.* 2007 Nov; 55:777-80.
- Moses AO, Munir'deen AI, Adegboyega FA and Abdulgafar JA. A study of serological evidence of prior chlamydia trachomatis infection in patients with ectopic pregnancy in Ilorin, Nigeria. *European Journal of Scientific Research.* 2007; 16(2):461-466.
- Jeremiah I, Okike O and Akanni C. The Prevalence of Serum Immunoglobulin G Antibody to Chlamydia Trachomatis in Subfertile Women Presenting at the University of Port Harcourt Teaching Hospital, Nigeria. *Int J Biomed Sci.* 2011; 7(2): 120–124.
- Mawak JD, Dashe N, Agabi YA and Panshak BW. Prevalence of Genital Chlamydia Trachomatis Infection among Gynaecology Clinic Attendees in Jos, Nigeria. *Shiraz E-Medical J.* 2011; 12:2

17. Okoror LE, Omilabu SA, Fadojutimi J and Nsongkhai V. Seroprevalence survey of Chlamydia in patients attending pre and postnatal clinic at the College of Medicine of the University of Lagos, Nigeria. Book of Abstract of the 24th Annual Conference of The Nigerian Society for Microbiology. 2000.
18. Arinze AU, Onyebuchi NV and Jeremiah I. Genital chlamydia trachomatis infection among female undergraduate students of University of Port Harcourt, Nigeria. Niger Med J. 2014; 55(1): 9–13
19. Nwanguma B, Kalu I and Ezeanyika L. Seroprevalence of anti-chlamydia trachomatis IgA antibody in a Nigeria population: Diagnostic significance and implication for the heterosexual transmission of HIV. The Internet Journal of Infectious Disease. 2008;7(2): 1-6
20. Ikeme AC, Ezegwui HU, Ikeako LC, Agbata I and Agbata E. Seroprevalence of Chlamydia trachomatis in Enugu, Nigeria. Niger J Clin Pract. 2011;14:176–80
21. Cuffini C, Bottiglieri M, Kigwen X, Alonso C, Deimundo S, Isa M, Cannistraci R, Gonzalez S and Farinati A. Molecular Epidemiology of genital Chlamydia trachomatis infection in asymptomatic adolescent young people. J Microbiol Research 2012; 2(4):114-117.
22. Alarape AI. Condom Use Among Students: The Influence of Condom Self-Efficacy, Social Norms and Affective Attitude Towards Condom. Journal of social sciences 2008; 17(3):237-241
23. Dean AG, Dean JA, Coulombier D, Buron AH, Brendel KA, Smith DC, Dicker R, Sullivan KM and Fagan RF. A word-processing, database and statistical program for public health on IBM-compatible microcomputers. Centers for Disease Control and Prevention, Atlanta, GA, USA. Epi. Info. Version 6.04 1996.
24. Theunissen JJ, Minderhoud-Bassie W, Wagenvoort JH, Stolz E, Michel MF and Huikeshoven FJM. Chlamydia trachomatis-specific antibodies in patients with pelvic inflammatory disease: comparison with isolation in tissue culture or detection with polymerase chain reaction. Genitourin Med 1994; 30:304-307
25. Fresse AS, Sueur JM and Hamdad F. Diagnosis and follow-up of genital chlamydial infection by direct methods and by detection of serum IgG, IgA and secretory IgA. Indian J. Med. Microbiol. 2010; 28(4):326-331
26. Van den Broek IVF, Land JA, Van Bergen JEM, Morré SA and Van der Sande MAB. Chlamydia trachomatis Antibody Testing in Vaginal Mucosal Material versus Blood Samples of Women Attending a Fertility Clinic and an STI Clinic. Obstet Gynecol Int. 2014; 2014: 601932. doi: 10.1155/2014/601932
27. Mazzoli S, Cai T, Rupealta V, Gavazzi A, Castricchi Pagliai R, Mondaini N and Bartoletti R. Interleukin 8 and anti-Chlamydia trachomatis mucosal IgA as urogenital immunologic markers in patients with C. trachomatis prostatic infection. Eur Urol 2007; 51:1385-93
28. Hartog JE, Land JA, Stassen FR, Kessels AG and Bruggema CA. Serological markers of persistent C. trachomatis infections in women with tubal factor subfertility. Human Reproduction. 2005;20 (4) :986–990.