

ORIGINAL RESEARCH ARTICLE

Incidence, trends and risk factors for perineal injuries of low-risk pregnant women: Experience from a midwife run obstetric unit, South Africa

DOI: 10.29063/ajrh2021/v25i4.6

Akm M. Hoque^{1*}, Muhammad E. Hoque² and Guido V. Hal³

Medical Manager, Kwadabeka Community Health Centre, KwaZulu-Natal, South Africa¹; Management College of Southern Africa Durban, South Africa²; Social Epidemiology and Health Policy, University of Antwerp Antwerp, Belgium³

***For Correspondence:** Email: mhoque75@gmail.com Phone: +27 31 7143704

Abstract

Pregnant women experience perineal injuries during childbirth. The objectives of this cross-sectional retrospective study were to estimate the incidence, trends, and risk factors for perineal injuries of women who had childbirths from January 2013 to December 2017. We used logistic regression to identify risk factors for all injuries, episiotomy, and obstetric anal sphincter injury (OASI) measured by odds ratios (OR). A total of 5547 women showed gradual decreases of episiotomy from 17.6% in 2013 to 7.6% in 2017 ($p < 0.05$). Perineal injuries were reduced from 33.3% in 2013 to 28.9% in 2017 ($p < 0.05$). The risk factor for any perineal injury were younger ages, term pregnancy, and nil parity ($p < 0.05$). Advanced gestational age, nil parity, and previous vaginal births were risk factors for episiotomy. However, birth weight of baby was significantly associated with OASI. Episiotomy and overall perineal injury rates were commendable. Training to midwives is needed to improve perineal care and maintain good practices during delivery. (*Afr J Reprod Health* 2021; 25[4]: 52-62).

Keywords: Episiotomy, labour management, obstetric anal sphincter injury

Résumé

Les femmes enceintes subissent des blessures périnéales lors des accouchements. Les objectifs de cette étude rétrospective transversale étaient d'estimer l'incidence, les tendances et les facteurs de risque de blessures périnéales chez les femmes qui ont accouché de janvier 2013 à décembre 2017. Nous avons utilisé la régression logistique pour identifier les facteurs de risque pour toutes les blessures, l'épisiotomie et l'obstétrique. Lésion du sphincter anal (OASI) mesurée par les rapports de cotes (OR). Au total, 5 547 femmes ont présenté une diminution progressive de l'épisiotomie de 17,6 % en 2013 à 7,6 % en 2017 ($p < 0,05$). Les blessures périnéales ont été réduites de 33,3% en 2013 à 28,9% en 2017 ($p < 0,05$). Le facteur de risque de toute lésion périnéale était un âge plus jeune, une grossesse à terme et une parité nulle ($p < 0,05$). L'âge gestationnel avancé, la parité nulle et les accouchements vaginaux antérieurs étaient des facteurs de risque d'épisiotomie. Cependant, le poids de naissance du bébé était significativement associé à l'OASI. Les taux d'épisiotomie et de lésions périnéales globales étaient louables. La formation des sages-femmes est nécessaire pour améliorer les soins périnéaux et maintenir les bonnes pratiques lors de l'accouchement. (*Afr J Reprod Health* 2021; 25[4]: 52-62).

Mots-clés: Épisiotomie, gestion du travail, lésion obstétricale du sphincter anal

Introduction

Pregnant women experience varying degrees of perineal injuries during childbirth. Perineal injuries are classified as (i) first degree: when the injury involves perineal skin, (ii) second degree: perineal muscles and skin are involved, (iii) third degree: injury involving anal sphincter complex, and (iv) fourth degree: involving the anal sphincter complex, epithelium, and rectal mucosa^{1,2}. The third- and

fourth-degree perineal injuries are considered major or extensive and are collectively known as obstetric anal sphincter injuries (OASI). The overall rates of OASI are reported with great variations from different reports between 0.7% and 10.2% (Norway, Denmark, Sweden, Iceland, USA, and UK) at different periods³⁻⁷. A recent study of a systemic review of meta-analysis on "Birth-Related Trauma in Low-and middle-Income Countries (LMIC)" reported that the overall episiotomy, second degree

injury, and OASI rates were 46%, 24%, and 1.4% respectively⁸. However, this report excludes first degree injuries as a result of incomplete data from the published reports. The study further identifies other limitations such as definition, reporting, and outcomes of perineal injuries, the use of episiotomy (routine versus selective) to prevent OASI and or to facilitate childbirths, and suggestions made for improvement on monitoring and reporting of perineal injuries from different health facilities of LMIC⁸. High incidence of all perineal injuries was reported between 70 to 85% from high income countries where the monitoring and reporting systems are of high quality and included all injuries^{9, 10}. A study from Sweden reported that perineal injuries are lower among planned home childbirths than hospital births¹¹. Higher incidences of overall perineal injuries were reported from England (90.4%) and Iran (84.3%)¹²⁻¹³. The lowest incidence of 64% perineal injury was reported from Brazil among low-risk pregnancies¹⁴.

It is reported that perineal injuries are related to pelvic floor disorders even after 10 years of delivery¹⁵. The incidence of episiotomy and OASI are also considered as obstetric care indicators^{12,16-17}. Detection and repair of extensive perineal injuries involving anal sphincter are thus important for maternity care. Therefore, it is universally recommended that the perineum is supported during the time of delivery of the foetal presenting part that causes stretching of the perineum as standard obstetric care and is found to prevent injuries^{18,19}. There are controversies on manual support to prevent perineal injuries. For example, a report from a meta-analysis of randomised controlled trials (RCTs) finds no protective effect but non-randomised studies find significant protection for OASI²⁰. Studies report from Norway that training of doctors and midwives on the traditional method of manual support with significant reductions on the incidences of OASI, overall spontaneous and operative vaginal deliveries^{21,22}. A randomised control trial on primiparous women from Brazil reports that the left lateral position during childbirth resulted in low risk of OASI²³. A similar reduction of perineal injuries including

episiotomy is found when women had childbirth on left lateral position compared to lithotomy position²⁴. Therefore, the left lateral position of women at the time of childbirth is considered protective for perineal injuries.

The known maternal risk factors for perineal injuries are age, parity, precipitated labour and very narrow introitus (foetal passage) lead to cephalo-pelvic disproportion (CPD) and foetal factors such as large foetus, occipito-posterior position of the vertex (foetal head), and or malpresentation. The known obstetric factors are uncontrolled or precipitated delivery (labour), assisted deliveries, episiotomy, vacuum extraction, and extended episiotomy in emergency lead to perineal injuries²⁵. A study from USA hospitals in 2011 on “third- and fourth-degree perineal tears prevalence and risk factors” reported that occipito-posterior position being the presenting part, parity and excessive birth weight of the new-born) were significant predictors of the cause of OASI²⁶. Another study on “risk factors for OASI during vaginal delivery from a referral hospital” in Cape Town, South Africa (SA), identified primipara, assisted childbirths (use of forceps and vacuums), malpresentation, mothers negative HIV status, and shoulder dystocia were significantly associated with perineal injuries²⁷. In that hospital, midwives only conducted uncomplicated childbirths while complicated births were assisted and conducted by medical professionals under Obstetrician’s supervision and used mediolateral episiotomies when necessary. Similarly, a report from two regional hospitals of Durban in SA found a rate of 16.2% spontaneous perineal injuries²⁸. The same study found race, the time required for childbirths, and the use of epidural analgesia were significantly associated with perineal injuries. Episiotomy was found to be a protective factor for OASI²⁸. There is limited information from low-income countries and more so from the midwife obstetric unit (MOU) where women give births. Therefore, it is important to understand the magnitude, classification, risk factors of perineal injuries from different settings. The objectives of this study are to estimate the incidence, trends, and risk factors for perineal injuries of women who gave childbirths at a MOU.

Methods

Study design

A cross-sectional retrospective study was undertaken to target all women who had spontaneous singleton vaginal childbirths at Kwadabeka community health center (KCHC) from January 2013 to December 2017.

Study setting and data collection

The setting of the study has been explained elsewhere as this was part of a comprehensive study that investigated the problems and outcomes of pregnant women experienced during the time of delivery²⁸. However, the study was undertaken at KCHC, a Primary Health Care (PHC) facility in Durban, SA, for the residence of Kwadabeka and Clermont communities with over 150,000. The maternity services at this MOU are available 24 hours a day and are run by trained midwives using SA National protocol²⁹.

Care and management of perineum during the second stage of labor²⁹

The second stage of labour is defined when the cervix is fully dilated and ends with the delivery of the baby. Usually, two hours are allowed for the foetal head or the presenting part to descend onto the pelvic floor if there is no foetal distress and CPD. The bladder is emptied using a catheter, if necessary, as usual practice for easy descent of the presenting part of the foetus. Delivery of the foetus is usually undertaken in lithotomy position as a routine practice at the facility. Efforts to bear down the foetus by the mothers are only encouraged when the foetal head starts to distend in the perineum with uterine contractions and the woman has an urge to push. When the woman is ready to bear down the baby, the woman is encouraged to bearing down only during contractions of the uterus (experience of pain by the mother). To protect the perineum, midwives use a perineal guard when the foetal head crowns. An episiotomy is considered and undertaken on selective cases such as thick or rigid perineum that seems to prevent delivery and may prolong the second stage of labour. The other maternal and foetal conditions for episiotomy are

signs of foetal distress in the second stage of labour, breech delivery, history of previous third- or fourth degree perineal injuries, or preterm delivery where the perineum is tight. A standard right mediolateral episiotomy is undertaken using local anaesthetic. However, there is no induction of labour and instrumental deliveries using forceps or vacuum extractor at this MOU by the midwives. A rectal examination after suturing the episiotomy or second-degree injury is performed by the delivering midwife to check for any stitches placed in the rectum.

Referral criteria of pregnant women from KCHC to hospitals during labour²⁹

Pregnant women attended KCHC in labour with the following conditions were referred to hospitals: primipara women aged ≥ 37 years, grand multiparity (parity ≥ 5), had previous caesarean section or surgery of the uterus, cervix, vagina, bladder or pelvic floor, previous postpartum haemorrhage requiring blood transfusion, serious medical disorder (e.g. cardiac disease, current TB infection, currently symptomatic asthma, epilepsy), anaemia (Hb < 10 g/dL), hypertension ($\geq 140/90$ mmHg), multiple pregnancies, breech presentation or transverse lie, estimated foetal weight < 2 kg, rupture of the membranes before the onset of labour, maternal pyrexia ≥ 37.5 degrees Celsius, vulvovaginal blisters or ulcers, extensive vulvovaginal warts that may obstruct delivery, antepartum haemorrhage, suspected foetal distress, thick meconium staining liquor, offensive liquor, cord prolapse, prolonged latent phase (≥ 8 hours) of labour, poor progress in the active phase (first stage) of labour (> 8 hours) and prolonged second stage of labour (> 2 hours).

Definition of terms

APGAR score stands for "Appearance, Pulse, Grimace, Activity, and Respiration" for the newborn babies in 1 and 5 minutes. Five indicators were used to check the health of the baby. Each indicator was scored on a scale of 0 to 2, with 2 being the best score. Preterm or premature babies were defined when babies were born < 37 weeks of gestation, or they can be small for their gestational age (37 weeks of gestation but baby weight < 2500

grams). Preterm delivery was considered when mothers delivered a baby between 28 weeks and 36 weeks of gestational age and the baby weights above 1000g. The “term delivery” was considered between 37 and 41 weeks of gestation. Any delivery that occurred at 42 completed weeks or afterward was considered as “post-term delivery”.

Data analysis

We entered data into Microsoft Excel for Windows and imported it into Statistical Package for Social Sciences (IBM SPSS) version 22.0 software for coding and analysis. We analysed the following variables: (a) maternal factors: age in years, parity (nil, 1-4 & ≥ 5), and previous vaginal birth, antenatal care history; (b) obstetric factors: gestational age in weeks, episiotomy undertaken and (c) foetal factors: weight of the new-born (baby) in kilogram (Kg), gender of the new-born (male or female) and APGAR score at 1 and 5 minutes. Primary outcome measures of the current study were perineal injuries first categorized into a) induced injury (episiotomy) and b) spontaneous injuries. Spontaneous injuries were further categorized as i) first ii) second iii) third and iv) fourth degree injuries. The demographic, baseline dependent, and outcome variables of women were summarized using descriptive summary measures: expressed as mean with standard deviation for continuous variables. We used percent for categorical variables. Cross-table analysis of independent and dependent variables was undertaken using Chi-square test (X^2) to identify the factors significantly associated with outcome variables. We used binary logistic regression analysis to determine possible predictors for outcome variables (separately for total, episiotomy, and OASI) and the results were expressed with adjusted odds ratios (OR) with corresponding two-sided 95% confidence intervals (95% CI) and associated p-values. P-values <0.05 were considered significant.

Results

A total of 5547 pregnant women had delivered singleton babies during 2013-2017 and thus formed our study sample. The mean age was 24.67 (SD= 5.89) years ranging from 13 to 47 years. Most of them (60%) belonged to the age group 20-29 years

(Table 1). Nearly all of them were at term gestation (97%), the majority (73.1%) had parity between 1 and 4, previous vaginal deliveries (73.3%), and received antenatal care (93.6%). The low-birth-weight delivery rate (≤ 2.5 kg) was 7.8% and most of the delivered babies (92.2%) had birth weight between 2.5 to 4.0 Kg. APGAR scores (over ≥ 7) of the babies in 1 and 5 minutes were 92.3% and 96.2% respectively.

The summary of all five years birth data showed (Table 1) that more than one fifth (21.1%) of the pregnant women had spontaneous perineal injuries while 11.3% had episiotomy making a total of 32.4%. Among spontaneous perineal injuries, the incidences of first- and second-degree injuries were 17.6% and 3.3% respectively. Only a few had third degree (0.2%) while none had fourth-degree perineal injuries. Those who had undergone episiotomy did not have further third- or fourth-degree injuries. Table 1 also depicted the cross-table analysis with Chi-Square (X^2) and p-values. There was a significantly higher (32.5%) rate of spontaneous perineal injuries among teenagers ($p<0.01$) compared to older women. A significantly higher rate (24.2%) of perineal injury was found among those women who had term pregnancy (gestational age ≥ 37 weeks)) compared to preterm (12.6%) ($p=0.002$). A higher rate (25.8%) of perineal injury was found among those women that delivered babies weighing between 3-3.49 kg compared to lower birth weight categories ($p<0.05$).

Figure 1 showed the trends of perineal injury rates over the study period. At the base year (2013) the total (all types) injury rate was higher of 33.3% and was found to decrease significantly to 28.9% ($p<0.05$) in 2017. The overall reduction of all perineal injuries was 13%. The episiotomy rates were also decreased from 17.6% in 2013 to 7.6% in 2017 ($p<0.05$) with a reduction of 57%. However, the spontaneous perineal injuries were increased significantly from 15.7% in 2013 to 21.3% ($p<0.05$) in 2017 with the highest rate of 25.2% in 2016. Binary logistic output (Table 2) on all perineal injuries showed that the younger pregnant women had higher risk of perineal injuries. We found that teenage age (< 20 years) and ages between 20-29 years were 6 ($p<0.05$), and 3.8 times ($p<0.05$) respectively more likely to have perineal injuries than the older women.

Table 1: Baseline variables with cross-table analysis with outcome variables of the study population

Variables	Frequency	%	Spontaneous Injury (%)	P-value	Episiotomy (%)	P-value	OASI (%)	P value
Age (n=5542)								
<20 years	1068	19.2	5.5	0.000	4.3	0.000	0.1	0.010
20-29 years	3332	59.9	14.7		6.1		0.1	
30- 39 years	1082	19.4	3.5		0.9		0.0	
>40 years	51	0.9	0.0		0.1		0.0	
Gestation age (n= 5508)								
Term (37 -40 weeks)	5338	96.9	23.4	0.001	11.2	0.000	0.2	0.298
Preterm (\leq 36 weeks)	170	3.1	0.4		0.1		0.0	
Parity (n= 5471)								
Nil parity	1386	25.3	7.6	0.000	5.6	0.000	0.1	0.845
1-4 parity	3996	73.1	16.1		5.6		0.2	
>5 parity	89	1.6	0.2		0.1		0.0	
Previous vaginal deliveries (n=5547)								
Yes	4073	73.3	6.1	0.449	3.6	0.001	0.1	0.919
No	1474	26.7	17.7		7.7		0.2	
Antenatal booking (n=5547)								
Yes	5181	93.6	1.3	0.011	0.3	0.000	0.0	0.367
No	357	6.4	22.6		10.9		0.2	
Sex of baby (n=5500)								
Male	2759	49.6	12.4	0.014	5.8	0.043	0.1	0.999
Female	2741	49.4	11.4		5.4		0.1	
Birth weight (n= 4874)								
< 3 Kgs	1748	31.4	6.2	0.00	3.3	0.189	0.0	0.000
3- 3.99 Kgs	3711	66.8	17.3		7.8		0.1	
\geq 4.00 Kgs	99	1.8	0.4		0.1		0.1	
Length (n=5210)								
\leq 50 cm	4167	80	19.7	0.542	9.3	0.120		
\geq 51 cm	1043	20	4.8		2.0			
APGAR score (n=5547)								
<7 in 1 minutes	425	7.7	0.8	0.003	0.7	0.162	0.0	0.440
\geq 7 in 1 minutes	5122	92.3	23.3		10.7		0.2	
< 7 in 5 min	212	3.8	0.0	0.001	0.1	0.462	0.0	0.740
\geq 7 in 5 min	5351	96.2	24.1		11.4		0.2	
Perineal injuries (n= 5545)								
Intact perineum	3747	67.6						
1-degree injury	978	17.6						
2-degree injury	184	3.3						
3-degree injury	11	0.2						
Episiotomy	625	11.3						

Table 2: Logistic regression output of all perineal injuries

Variables	Sig.	Odds (OR)	Ratio	95% C.I. for OR	
				Lower	Upper
Age coded	.000				
Age < 20 years	.001	5.927		2.084	16.861
Age 20-29 years	.011	3.866		1.369	10.920
Age 30-39 years	.085	2.506		.882	7.120
Gestational age \geq 37 weeks	.003	2.046		1.286	3.256
Parity	.000				
Parity nil	.000	3.919		2.021	7.600
Parity 1-2	.072	1.817		.948	3.484
Birth weight of the baby	.000				
Birth weight < 3 kgs	.868	.960		.592	1.555
Birth weight 3-3.99 Kgs	.242	1.327		.826	2.130
ANC booking (Yes)	.008	.689		.524	.905
Constant	.000	.024			




Figure 1: Trends of different types of perineal injuries from 2013 to 2017 at KCHC

Table 3: Logistic regression output for episiotomy

Variables	Sig.	Odds (OR)	Ratio95% C.I. for OR	
			Lower	Upper
Age coded	.000			
Age < 20 years	.176	2.303	.689	7.701
Age 20-29 years	.711	1.254	.378	4.159
Age 30-39 years	.534	.679	.200	2.301
Gestational age \geq 37 weeks	.000	6.226	2.250	17.232
Parity coded	.000			
Parity Nil	.024	3.973	1.202	13.130
Parity 1-4	.448	1.583	.484	5.180
Had previous vaginal delivery	.016	1.262	1.045	1.525
Had antenatal booking	.002	.457	.279	.749
APGAR scores > 7 after 5 minutes of delivery	.078	1.397	.963	2.026
Constant	.000	.007		

Table 4: Logistic regression output for OASI

Variables	Sig.	Odds ratio (OR)	95% C.I. for OR	
			Lower	Upper
Birth weight	.000			
Birth weight < 3 Kgs	.000	.14	.011	.142
Birth weight 3-3.99 Kgs	.000	.55	.133	.223
Gestational age < 37 weeks	.066	.115	.011	1.157
Constant	.997	.000		

Primipara women and women with gestational age at term (≥ 37 weeks) were 3.9 ($p < 0.05$) and 2 times ($p < 0.05$) more likely to have perineal injuries than their counterparts. Primipara women were almost 4 times ($OR = 3.97$, $p < 0.05$) more likely to have undergone episiotomy (Table 3) compared to

multipara Pregnant women at term pregnancy (> 37 weeks) were 6 times ($OR = 6.22$, $p < .001$) more likely to have episiotomy than those had preterm childbirths. Similarly, women who had a previous vaginal delivery were 1.2 ($OR = 1.2$, $p < 0.05$) times more likely to have an episiotomy than those that

did not have previous vaginal deliveries. On the other hand, pregnant women who attended antenatal care were 55% less likely to have undergone episiotomy (OR=0.457, $p<0.05$) than those who did not. Lower birth weights of the babies were found to be a protective factor for OASI. Result (Table 4) showed that the birth weights of the babies < 3.0 Kg and between 3.0-3.99 Kg were 86% (OR=0.14, $p<0.05$) and 45% (OR=0.55, $p <0.05$) less likely respectively to have OASI compared to birth weight of ≥ 4 kgs. No demographic, obstetrics variables, and episiotomy were associated with OASI.

Discussion

This study collected data from a large number of vaginal deliveries from low-risk pregnant women between 2013 and 2017 and estimated the incidences and risk factors for different degrees of perineal injuries. Firstly, we found trends of all injuries over 5 years period (spontaneous, episiotomy, and total) and secondly incidence of total injuries of different degrees and their risk factors. We found a decreasing trend of episiotomy from a higher rate of 17.6% in 2013 to a lower rate of 7.6% in 2017, a reduction of 57%. The episiotomy rate in 2017 had reached the rate recommended by WHO and is similar to other findings from Africa^{16,30,31}. However, this trend must be seen with the increasing trend of spontaneous perineal injuries over the same period. There was an increase in spontaneous injury (21.3%) in 2017 compared to the rate of 15.7% in 2013 (Figure 1). It can be argued that it was due to the reduction of episiotomy, there was an increase of spontaneous perineal injuries. However, there was a significant reduction (13%) of all perineal injuries from 2013 to 2017. These decreasing trends of total perineal injuries including episiotomy indicated that there were constant efforts to avoid unnecessary use of surgical intervention (episiotomy) and to prevent any spontaneous injuries for a physiological process of vaginal delivery. The reduction rates should be seen positively as midwives are constantly striving towards better services for pregnant women and minimizing preventable perineal injuries at this MOU. However, good practices of perineal care during childbirth such as the position of women

(lithotomy, standing, sitting, left lateral, etc.), perineal support, bearing down of babies all contributed to the reduction of episiotomy and overall perineal injuries.

Five years summary data showed that more than one fifth (21.1%) of these women had spontaneous and 11.3% had induced (episiotomy) perineal injuries. The incidence of spontaneous perineal injuries among the low-risk pregnant women appeared higher than other reports from similar resource-constrained settings^{11,28,32}. The rate was higher than the rate found from a hospital delivery in Durban (16.2%), SA²⁸. This could be due to the support that midwives received from doctors in the hospital. The incidence of perineal injuries was also found to be higher among black South African pregnant women than the other races in SA and elsewhere^{28,33}. The rates of the second degree and OASI were minimum in our setting compared to other studies in hospital settings in SA and abroad^{14,27-28}. Regional hospitals in SA conduct deliveries of complicated pregnancies referred from MOUs and district hospitals thus a higher rate of OASI is likely. The incidences of perineal injuries were found to markedly vary between different study settings with higher rates in hospitals compared to lower in community settings and those were found from Sweden and Nicaragua^{11,34}. The first degree perineal injury is considered minor and it was 18% in our study. We found a low incidence of the second degree and OASI (3.3% and 0.2% respectively). The spontaneous second-degree perineal injuries were much lower than the reported rate from a meta-analysis (23%)⁸. However, the rate of second-degree perineal injury in our study is comparable with other findings from Nicaragua (2.7%), Pakistan (3.2%), and Bangladesh (1.1%)³⁴⁻³⁶. The second degree perineal injury though considered a minor injury still needed special attention as it affects the perineal muscles. Though muscular injury is classified as a second-degree injury it is equal to and often becomes worse than a routine episiotomy. However, if the injury involves the levator ani muscle it leads to pelvic floor disorders in later life¹⁰.

Risk factors for spontaneous perineal injuries were well documented in previous reports. Teenagers (age < 20 years) and ages between 20-29 years), nulliparous (parity nil) pregnant women

showed strong risk factors for spontaneous perineal injuries in our study which were well recognized in earlier reports^{24,37}. In our study, term pregnancy had twice the chance of having spontaneous injuries. Not many studies looked at gestational age as a risk factor for perineal injury. The case-control study from Cape Town tertiary hospital (SA) looked into it and reported that gestational age was not a risk factor²⁷. As the gestational age increases, the foetus grows bigger and gains weight. These two factors (higher gestational age and baby weight) are interrelated. We found both factors were indeed risk factors for perineal injuries like other study²⁶.

The episiotomy rate in our study was lower compared to other reports from Ethiopia where they found a rate of 35%, France (national average of 14.1% for all non-operative vaginal deliveries), and Vietnam (15.1%)³⁸⁻⁴⁰. However, the incidence of episiotomy is similar to the rate (10%) recommended by WHO when it is undertaken for selected cases¹⁶. However, the episiotomy rate in our study is higher than the rates reported from Brazil (8%) for low-risk pregnancies delivered at a hospital referred from PHC clinics and in Nigeria (9.3%)^{14,30}. The possible reason for this low incidence of episiotomy in our study was because episiotomy was undertaken when it was indicated, and also to the fact that the low-risk pregnant women delivered at this MOU²⁹. Routine episiotomy in reducing severe or major perineal injury (third or fourth degrees) became a controversial issue in modern obstetric practice. There was a systematic review report that supported that standard episiotomy (at mediolateral position) was found to reduce severe forms of perineal injury (third- and fourth-degree injuries)³¹. Similarly, episiotomy was found with an association of reducing perineal injuries in hospital deliveries from Durban, SA²⁸. On the contrary, other reports from Cape Town and the Cochrane study found that there was no association to protect major perineal injuries using routine episiotomy^{27,41}. We also did not find any association between episiotomy and OASI. A report from Australia indicated that episiotomy was associated with minimising major injuries when assisted vaginal delivery was conducted using forceps⁴². In our set up there was no forceps assisted deliveries undertaken. The factors found associated with undertaking episiotomy in our study were

gestational age ≥ 37 weeks, primipara, and previous vaginal deliveries which are similar to other reports from Africa and elsewhere^{38,39,43}. Pregnant women who had antenatal care during pregnancy were found to be protective (55%) against episiotomy. There is no report which highlighted that antenatal care during the antenatal period could reduce perineal injuries, especially episiotomy. However, it was reported that antenatal education prepares pregnant women for delivery and found to impact positively on wound-healing and compliance with wound care⁴⁴.

The OASI in our study was low of 0.2%. This is lower than the rates found in Brazil (0.75% for low-risk pregnancies) and Mexico (0.8% for all vaginal deliveries)^{14,32}. The reason for the low rate of OASI in our study could well be due to low-risk pregnant women who delivered in our facility. The misdiagnosis and underreporting of major perineal injuries cannot be ruled out as suggested by others³². However, underreporting is less likely as all OASI cases were needed to be referred to a hospital for surgical intervention like suturing of an anal sphincter. Several studies suggested that the heavier birth weight of babies at delivery was associated with severe perineal injuries. Our study found a similar trend of lower birth weight (<4Kg) was protective for OASI^{11,32,33}. The earlier studies from Brazil and Mexico reported that women with younger age (teenage women) had 1.3 and 2.9 times respectively more likely to have OASI^{14,32}. This was not the case in our study. In an earlier report, it was found that the negative impact of OASI in subsequent pregnancies was five-fold to have severe perineal injuries thus strategies should be considered essential to prevent OASI⁴⁹.

Strengths and limitations

The strength of this study was its fairly large homogenous type of sample. The limitations of this study were its retrospective nature and review of records limited the study variables and made us reliant on the quality of data recorded. Finally, more prospective studies are necessary, in order to assess more risk factors associated with mild and severe perineal injuries. It is important to identify women who are at risk of OASI during childbirth, in order to minimize the risks of perineal injury during this

period. Midwives from PHC facilities need to have advanced knowledge of pelvic and perineal anatomy, so as to prevent injuries during childbirth.

Conclusion

This large sample of women from a MOU had shown declining trends of perineal injuries. Risk factors identified for these injuries are similar to those previously reported in other studies carried out in different settings. Identification of those parturient women who are at higher risk groups may result in timely and appropriate interventions that minimize perineal injuries and complications thus preventing postpartum and the long-term sequelae that may develop later in life. Further studies are recommended to identify the effect of antenatal care on perineal health at the time of delivery and to monitor the trends of perineal injuries.

Acknowledgements

We thank all practicing midwives of KCHC working in labour ward for recording study variables in the birth register, Maariyah Hoque who captured data in Microsoft Excel, and for coding and assisting in the analysis of data in SPSS for the report.

Ethical approval

We obtained ethical approval from Umgungundlovu Health Ethics Research Board (Reference no. UHERB 015/2020). We sought permission from the management of KCHC for utilizing the delivery register to conduct the study. We did not use the identification of patients or staff during the analysis and presentation of the results.

Funding

None

Conflict of interest

There is no competing interest of authors to this report and study.

Contributions of authors

AMH – Conceptualisation, study design, monitoring and participation in data collection,

collation and analysis, preparation, and finalization of the manuscript.

MEH– Conceptualisation, data analysis, editing, and finalization of the manuscript

GVH – Conceptualisation, editing, and finalization of the manuscript. All authors have read and approved the manuscript.

References

1. World Health Organization. International Classification of Diseases (ICD). Geneva (CG): WHO; 2015. Available at: <http://www.who.int/classifications/icd/en>.
2. Fernando RJ, Williams AA and Adams EJ. The Management of Third and Fourth Degree Perineal Tears. RCOG Green top Guidelines No 29. London: RCOG, 2007
3. Edqvist M, Blix E, Hegaard HK, Ólafsdóttir OA, Hildingsson I, Ingversen K, Mollberg M and Lindgren H. Perineal injuries and birth positions among 2992 women with a low risk pregnancy who opted for a homebirth. BMC Pregnancy and Childbirth 2016; 16:196.
4. Minaglia SM, Ozel B, Gatto NM, Korst L, Mishell DRJ and Miller DA. Decreased rate of obstetrical anal sphincter laceration is associated with change in obstetric practice. Int Urogynecol J Pelvic Floor Dysfunct 2007;18:1399–404.
5. LaCross A, Groff M and Smaldone A. Obstetric anal sphincter injury and anal incontinence following vaginal birth: a systematic review and meta-analysis. J Midwifery Womens Health. 2015;60:37–47
6. Raisanen S, Vehviläinen-Julkunen K, Gissler M and Heinonen S. High episiotomy rate protects from obstetric anal sphincter ruptures: a birth register-study on delivery intervention policies in Finland. Scand J Public Health. 2011; 39:457–63.
7. Valbo A, Gjessing L, Herzog C, Goderstad JM, Laine K and Valset AM. Anal sphincter tears at spontaneous delivery: a comparison of five hospitals in Norway. Acta Obstet Gynecol Scand. 2016; 95:941–947.
8. Aguiar M, Farley A, Hope L, Amin A, Shah P and anaseki-Holland S. Birth-Related Perineal Trauma in Low- and Middle-Income Countries: A Systematic Review and Meta-analysis. Matern Child Health J. 2019;23(8):1048-1070.
9. Rikard-Bell J, Iyer J and Rane A. Perineal outcome and the risk of pelvic floor dysfunction: a cohort study of primiparous women. Aust N Z J Obstet Gynaecol 2014; 54:371-376.
10. Webb S, Sherburn M and Ismail KMK. Managing perineal trauma after childbirth. BMJ 2014;25(349):6829-6829.
11. Lindgren HE, Radestad IJ, Christensson K and Hildingsson IM: Outcome of planned home births compared to hospital births in Sweden between 1992 and 2004. A population-based register study. Acta Obstet Gynecol Scand 2008; 87:751–759.

12. Smith L A, Price N, Simonite V and Burns EE. Incidence of and risk factors for perineal trauma: a prospective observational study. *BMC Pregnancy and Childbirth* 2013; 13: 59.
13. Abedzadeh-Kalahroudi M, Talebian A, Sadat Z and Mesdaghinia E. Perineal trauma: incidence and its risk factors *J Obstet Gynaecol* 2018; 39(2):1-6.
14. Peppe MV, Stefanello J, Infante BF, Kobayashi MT, Baraldi CO and Brito LGO. Perineal Trauma in a Low-risk Maternity with High Prevalence of Upright Position during the Second Stage of Labor. *Rev Bras Ginecol Obstet.* 2018; 40(7):379-383.
15. Handa VL, Blomquist JL, McDermott KC, Friedman S and Muñoz A. Pelvic floor disorders after vaginal birth: effect of episiotomy, perineal laceration, and operative birth. *Obstet Gynecol.* 2012;119:233- 9.
16. World Health Organization. Standards for improving quality of maternal and newborn care in health facilities. Geneva: World Health Organization, 2016
17. Australian Institute of Health and Welfare. Towards National Indicators of Safety and Quality in Health Care. Cat. no. HSE 75. Canberra: AIHW, 2009.
18. Grobler CJF and Cronje HS. Obstetrics in Southern Africa. 2003, Van Schaik, Hatfield, South Africa.
19. Mantzius J, Rudnicki M, Skovbjerg E, Maro E, Mrema D, Rasch V and Sørensen BL. Perineal trauma following vaginal delivery in a low-income area: A criterion-based audit. *Int. J. Nurs. Midwifery* 2020; 12(4): 113-119.
20. Bulchandani S, Watts E, Sucharitha A, Yates D and Ismail KM. Manual perineal support at the time of childbirth: a systematic review and meta-analysis. *BJOG* 2015;122:1157–1165
21. Hals E, Øian P, Pirhonen T, Gissler M, Hjelle S, Nilsen EB, Severinsen AM, Solsletten C, Hartgill T and Pirhonen J. A multicenter interventional program to reduce the incidence of anal sphincter tears. *Obstet Gynecol.* 2010;116(4):901-908.
22. Laine K, Skjeldestad FE, Sandvik L and Staff AC. Incidence of obstetric anal sphincter injuries after training to protect the perineum: cohort study. *BMJ Open* 2012;2:e001649.
23. Schirmer J, Fustinoni SM and Basile ALO. Perineal outcomes on the left lateral versus vertical semisitting birth positions: a randomized study. *Acta Paul Enferm* 2011;24(6):745-50.
24. Meyvis I, Van Rompaey B, Goormans K, Truijzen S, Lambers S, Mestdagh E and Mistiaen W. Maternal position and other variables: effects on perineal outcomes in 557 births. *Birth* 2012;39(2):115-20.
25. Dahlen HG, Ryan M, Homer CSE and Cooke M. An Australian prospective cohort study of risk factors for severe perineal trauma during childbirth. *Midwifery* 2007;23: 196-203.
26. Groutz A, Hasson J, Wengier A, Gold R, Skornick-Rapaport A, Lessing JB and Gordon D. Third- and fourth-degree perineal tears: prevalence and risk factors in the third millennium. *Am J Obstet Gynecol.* 2011; 204:347.1-4.
27. Juul L and Theron G B. Risk factors for third- and fourth-degree perineal tears during vaginal delivery. *Urogynaecologia.* 2011; 25:2.
28. Naidoo TD and Moodley J. Obstetric perineal injury: risk factors and prevalence in a resource-constrained setting. *Trop Doct.* 2015;45(4):252-4.
29. Department of Health. Guidelines for Maternity Care in South Africa: A Manual for Clinics, Community Health Centres and District Hospitals. 4th ed. Pretoria, South Africa: Department of Health; 2015.
30. Alayande BT, Amole IO and Olaoluva DA. Relative frequency and predictor of episiotomy in ogbomosho, Nigeria. *Internet J of Med Update.* 2012; 7:41-44.
31. Verghese TS, Champaneria R, Kapoor DS and Latthe PM. Obstetric anal sphincter injuries after episiotomy: Systematic review and meta-analysis. *Int Urogynecol J* 2016;27(10), 1459–1467.
32. Hirayama F, Koyanagi A, Mori R, Zhang J, Souza JP and Gu' Imezoglu AM. Prevalence and risk factors for third- and fourth-degree perineal lacerations during vaginal delivery: a multi-country study. *BJOG.* 2012; 119: 340–347.
33. Ampt AJ, Ford JB, Roberts CL and Morris JM. Trends in obstetric anal sphincter injuries and associated risk factors for vaginal singleton term births in New South Wales 2001–2009. *Aust NZ J Obstet Gynaecol* 2013; 53:9–16.
34. Garcia-Elorrio E, Aleman A, Cafferata ML, Colomar M and Tomasso GA multifaceted intervention to increase prophylactic oxytocin use during the third stage of labor and to reduce routine episiotomies in Nicaragua. *Int J Gynaecol Obstet* 2014;127(1):31–34.
35. Brohi ZP, Sadaf A, Zohra N and Perveen U. Frequency and severity of perineal tears in Countess Lady Duffrin Fund Hospital, Hyderabad. *J Pak Med Assoc.* 2012 Aug;62(8):803-6.
36. Ferdous J, Ahmed A, Dasgupta SK, Jahan M, Huda FA, Ronsmans C, Koblinsky M and Chowdhury ME. Occurrence and determinants of postpartum maternal morbidities and disabilities among women in Matlab, Bangladesh. *J Health Popul Nutr.* 2012;30(2):143-58.
37. Smith L A, Price N, Simonite V and Burns EE. Incidence of and risk factors for perineal trauma: A prospective observational study. *BMC Pregnancy and Childbirth.* 2013;13(1), 59-64.
38. Worku SA, Mitku YM and Getahun SA. Episiotomy practice and its associated factor among women who gave birth at public health institutions of akaki kaliti in Addis Ababa, Ethiopia. *Clinics Mother Child Health.* 16:318. doi:10.24105/2090-7214.16.318.
39. Goueslard K, Cottenet J, Roussot A, Clesse C, Sagot P and Quantin C. How did episiotomy rates change from 2007 to 2014? Population-based study in France. *BMC Pregnancy Childbirth* 2018; (18): 1747-58.
40. Trinh AT and Ampt KA. Episiotomy rate in vietnamese born women in Australia support for a change in obstetric practice in Vietnam. *Bull WHO.* 2013; 96:350-356.
41. Jiang H, Qian X, Carroli G and Garner P. Selective versus

- routine use of episiotomy for vaginal birth. The Cochrane Database of Systematic Reviews. 2017
42. Baghurst PA and Antoniou G. Risk Models for Benchmarking Severe Perineal Tears during Vaginal Childbirth: a Cross-sectional Study of Public Hospitals in South Australia, 2002–08. *Paediatric and perinatal epidemiology*. 2012;26(5):430-7.
43. Dannecker C, Hillemanns P, Strauss A, Hasbargen U, Hepp H and Anthuber C. Episiotomy and perineal tears presumed to be imminent: randomized controlled trial. *Acta Obstet Gynecol Scand*. 2014;83(4):364-8.
44. O’Kelly SM and Moore ZEH. Antenatal maternal education for improving postnatal perineal healing for women who have birthed in a hospital setting. Cochrane Database of Systematic Reviews 2017, Issue 12. Art. No.: CD012258.
45. Edozien LC, Gurol-Urganci I, Cromwell DA, Adams EJ, Richmond DH, Mahmood TA and van der Meulen JH. Impact of third- and fourth-degree perineal tears at first birth on subsequent pregnancy outcomes: a cohort study. *BJOG* 2014;121(13):1695–1703.