

ORIGINAL RESEARCH ARTICLE

Ricinus communis-linn (castor plant), male contraceptives and reproductive health of women

DOI: 10.29063/ajrh2021/v25i4.14

RT McNeil^{1*}, A Adebisin¹ and E Ekwere²

Department of Anatomy and Histology, Sefako Makgatho Health Services University, Pretoria, South Africa¹;
Department of Anatomy, University of Jos, Jos Nigeria²

*For Correspondence: Email: rosie.mcneil@smu.ac.za; Phone: 27 71 975 9167

Abstract

The medicinal use of castor plant is extensive. Castor oil is most commonly used as a laxative, and the leaves and seeds have been used to augment labour, promote lactation and to treat syphilis and leprosy. Its use for contraception is an ancient practice among the Rukuba women of Plateau State in Nigeria, who would chew 2-3 seeds for contraceptive coverage of one year. The acute and chronic spermatogenic effect of the minor seed variety of *ricinus communis-linn* is hereby reported. Thirty male rats weighing 200–250g were grouped (n = 5) and treated with the *n*-hexane extract of the seeds, each made up to 1ml with physiological saline, and given as a single dose, intraperitoneally. Control groups had 1 ml physiological saline. Semen was collected 72 h and 6 weeks after treatment and analyzed within one hour of collection. The seed extract suppressed spermatogenesis and sperm motility up to six weeks after treatment. But semen parameters reverted to normal values in the 7 weeks of treatment, showing primary spermatocytes and spermatids in the seminiferous tubules. Our results showed *ricinus communis-linn* to be a potent but reversible anti-spermatogenic agent with significant anti-motility properties. Moreover, being of plant origin, it is readily available and affordable. We hereby present *ricinus communis-linn* as a possible male contraceptive agent that can relieve women of their reproductive burden globally. (*Afr J Reprod Health* 2021; 25[4]: 135-141).

Keywords: *Ricinus Communis-Linn*, male contraception, reproductive health, women

Résumé

L'utilisation médicinale de la plante de ricin est étendue. L'huile de ricin est le plus souvent utilisée comme laxatif, et les feuilles et les graines ont été utilisées pour augmenter le travail, favoriser la lactation et traiter la syphilis et la lèpre. Son utilisation pour la contraception est une pratique ancienne chez les femmes Rukuba de l'État du Plateau au Nigeria, qui mâchaient 2-3 graines pour une couverture contraceptive d'un an. L'effet spermatogène aigu et chronique de la variété mineure de graines de *ricinus communis-linn* est rapporté ici. Trente rats mâles pesant 200 à 250 g ont été regroupés (n = 5) et traités avec l'extrait de *n*-hexane des graines, chacun complété à 1 ml de sérum physiologique et administré en une seule dose, par voie intrapéritonéale. Les groupes témoins avaient 1 ml de sérum physiologique. Le sperme a été collecté 72 h et 6 semaines après le traitement et analysé dans l'heure suivant le prélèvement. L'extrait de graines a supprimé la spermatogenèse et la motilité des spermatozoïdes jusqu'à six semaines après le traitement. Mais les paramètres du sperme sont revenus à des valeurs normales au cours des 7 semaines de traitement, montrant des spermatocytes et des spermatides primaires dans les tubules séminifères. Nos résultats ont montré que le *ricinus communis-linn* est un agent anti-spermatogène puissant mais réversible avec des propriétés anti-motilité significatives. De plus, étant d'origine végétale, il est facilement disponible et abordable. Nous présentons ici le *ricinus communis-linn* comme un agent contraceptif masculin possible qui peut soulager les femmes de leur fardeau reproductif à l'échelle mondiale. (*Afr J Reprod Health* 2021; 25[4]: 135-141).

Mots-clés: *Ricinus Communis-Linn*, contraception masculine, sante reproductive, femmes

Introduction

Family planning has been proven to improve maternal and child health, but the accompanying side effects present a major clinical problem¹⁻². While there are many successful female contraceptives, vasectomy and condoms are the most effective in males³. It has been said that all of

the contraceptive methods known, present with one or more problems¹. Methods currently used by females include injectables, intra uterine devices (IUDs), vaginal rings, patches, implants, condoms, oral pills, sterilization on the one hand, and lactational amenorrhea techniques, coitus interruptus, and awareness-based methods on the other hand⁴. WHO records of 2011 however, show

that 210 million pregnancies occur annually all over the world. Out of this, 80 million are unplanned, thus promoting unsafe abortion⁵ and avoidance of conventional contraceptives⁵. These range from weight gain and menstrual bleeding to cardiovascular complications and cancers⁶ and are found to be minimal with a reduction in the hormonal dosage². Irregular bleeding is the commonest reason for the discontinuation of problematic contraceptive methods⁴. Over the years, it has become certain that the use of current birth control methods presents with health complications that are severe enough to warrant clinical care and management of these side effects⁷. Because conventional female contraceptive agents often have severe health repercussions and since male anti-fertility agents are uncommon, some couples may be left with no contraceptive options. This calls for the need to find alternative methods that are less problematic to women of reproductive age. Even better still, a non-hormonal male contraceptive drug since most male contraceptives used in clinical trials consist of testosterone combined with other hormones. Unfortunately, the effect of long-term use is yet to be ascertained⁸. Furthermore, efforts are being made to find male contraceptive agents that have potent anti-spermatogenic effects but do not interfere with Leydig cell function. Spermatogenesis is a continuous process that begins at puberty and is regulated by pituitary testicular and adrenal hormones. Spermatocytes take approximately 72 days to mature. They travel through the seminiferous tubules to the epididymis, where they are stored prior to ejaculation².

Castor plant

Castor plant also called castor oil plant and castor beans belongs to the family of Eupobiaceae and is popular for its yield of castor oil. It is commonly a shrub of 8-12 feet but can be 20-30 feet (e"10m) high⁹ (USDA, 2016). Each tree bears both male and female flowers, which occur in clusters, upper (male) and lower (female) flowers.

The seed has three lobes, each lobe containing a seed, which is smooth and elliptical in shape¹⁰. They exhibit extreme variation in size and colour; they may be grey, brown, black, or brown/black mottled with a size of 0.8-1.8cm in

length, 0.4-1.2 cm in width, and oval or compressed in shape¹¹. They are 17 varieties of the seeds¹¹ which are classified into 3 varieties; variety major (or the large seed variety) variety intermedia (or medium seed variety) and various minor or small seed variety).

Many advances have been made in the molecular biology of castor seeds. Its seedlings contained more than 100 different polypeptides against which a complex of antiserum has been raised¹². Furthermore, the castor plant expressed auxin-related gene family, that regulates hormone signaling pathways in plants. These genes, therefore, regulate growth and stress response and have been associated with functional diversification in plants¹³. The seed contains about 20% shell and 80% kernel. Both consist of about 58-66% oil¹⁴. Trease and Evans recorded a fixed oil yield from the castor seed, of 55-60%. Other constituents include 20% proteins; ricin (toalbumin) albumin; enzymes (e.g. lipase, chymase); nuclealbumin; amino acids; globulins, glycoproteins, and alkaloids, steroids, and elements (magnesium, calcium, and manganese. Its toxicity is attributed to the protein content, ricin, constituting 5%^{15,11}.

Clinical trials with *Ricinus Communis-linn*

The use of castor seeds for contraception is widespread among the Rukuba tribal women of Bassa local government area in Plateau State, Nigeria¹⁶⁻¹⁸. In 1990, this property of castor seeds was thrown into the limelight when a female traditional doctor (called Hazija), communicated that 2-3 seeds protect the female from conception for a whole year¹⁹⁻²⁰. This information has led to an enormous amount of research work on the seed between 1990 until date. Prior to this, most work done on castor seed has been from the aspect of toxicity. More interestingly, some other researchers have reported on the antifertility property of the seed²¹⁻²². Furthermore, 10 women volunteers, oral administration of 3 seeds prevented contraception for 12 months with no sign of toxicity¹⁹.

Many plants and herbs have contraceptive properties^{19,23-24}. *Ricinus Communis-linn* showed an anti-ovulatory effect in Wistar rats²⁵ and in a reversible manner²⁵⁻³⁰. However, there is a dearth of data on its mechanism of action and antifertility

properties in male animal subjects. Moreover, this has not been investigated in men. Thus, the aim of this work is to investigate the anti spermatogenic properties of the N-Hexane extract of *ricinus communis-linn* in adult male Wistar rats.

Methods

Seed extraction

The seeds were extracted after the method of Okwuasaba *et al*³¹: Dried seeds (100 g) were ground into a fine powder using a mortar and pestle and soxhlet extracted for 72 h in 350 ml of *n*-hexane at 30°C. The extract was then concentrated in a water bath at 59.0 ± 1°C, dissolved in an appropriate volume of corn oil, and stored at 4°C.

Animal treatment

With ethical clearance, thirty male Wistar rats were raised to weigh between 200 and 250g and acclimatized for two weeks prior to the experiment. They were divided into treatment groups (n = 5). 10 mg/kg extract was administered to each test rat intraperitoneally through the anterior abdominal wall. Out of 10 test rats, 5 were euthanized after 72 h to observe the acute effect of the seed extract on semen parameters, and the other set (5), after 6 weeks in order to observe the chronic effect of the extract on semen quality. 10 control rats had equal volumes of physiological saline, 5 rats were sacrificed after 72 h and the other 5 after 6 weeks. All subjects were weighed weekly, fed with Pfizer pellets, and had water *ad libitum*. They were sacrificed in the 7th week of treatment using a chloroform chamber. The testes were exposed by an incision on the ventral surface of the scrotum, through which the caudal epididymis was excised, fenestrated, and placed in 1 ml of saline for semen analysis. A scientist that was blind to the experiment analyzed all semen samples within one hour of collection.

Results


As determined by microscopic analyses, the appearance, liquefaction, and consistency of all semen samples were normal. The extract suppressed semen parameters (mean count,

motility, and morphology) within 72 hours of treatment; there was a 50% reduction in the percentage mean normal forms (40%) when compared to the control subjects (80%). While the percentage mean motility was reduced to 56% in test rats, (control value was 88.0%). The percentage mean count was the most affected (49 (x10⁶). It was 99.5 (x10⁶) in the control group. Mean normal forms and motility improved over the next few weeks to 60 and 65 respectively but the mean count was 39 (x10⁶) in the 6th week, returning to values comparable with those of control rats only in the 8th week of treatment. Unlike Mean normal forms and Mean motility that showed significant improvement within six weeks, semen counts began to recover only at that time. This is attributed to the fact that the formation and maturation of spermatocytes take about 72 days³².

Discussion

In our previous work, we found castor seed to be a potent anti-ovulatory agent with oestrogen-like properties. In addition, cervical screening of women who had been on castor seeds for over five years yielded no side effects, as is common with conventional hormonal medicines. Female contraceptives are diverse, but male contraceptives tend to target semen parameters and are therefore specific in their function - to prevent pregnancy⁸. The seed of *ricinus communis-linn* is widely used around the world for traditional medicinal practices⁹. The seed extract suppressed the ovarian cycle of adult cyclic Sprague-Dawley rats⁶ and suppressed testicular function in a way similar to the anti-androgenic drug, cyproterone acetate⁷. Semen parameters determine male fertility¹⁰. Routine semen analyses assess multiple parameters including appearance, volume, liquefaction, concentration or viscosity, agglutination, hydrogen ion concentration, leucocyte count, sperm count, motility, viability and vitality, the morphology of spermatocytes, and anti-sperm antibodies¹¹.

Semen samples from the treatment groups showed azoospermia, with the percentage of normal morphologies being 40% (acute effect experiment), compared to 80% for the solvent control subjects. These values both improved to 60% and 86% respectively after six weeks of treatment.



A), fresh clusters of castor seed, B) and dried castor seeds, C)

Figure 1: The minor seed variety castor plant




shows the sizes of the seeds in centimeters; each seed is about 1.1cm long. The seeds were obtained from the Department of Pharmacology, University of Jos, Jos, Plateau State, Nigeria

Figure 2: The minor seed variety of castor plant (D) used in this study. Image E)


Why improvement in the control semen parameters? This may be because of relief from the stress of being experimented upon. The sympathetic nervous system and the adrenal cortisol regulate acute and chronic stress respectively. Unfortunately, these responses to stress, affect negatively, the reproductive system¹². Moreover, 5% of cases of infertility are due to emotional factors with neuroendocrinology and reproductive disruption¹³. Stress has been linked to poor semen quality; stress inhibits both the central control of reproduction and local testicular response (the hypothalamic–pituitary–gonadal axis). This

reduces testicular androgen production¹⁴. Reducing stress in the subjects may therefore alter semen values. Therefore, in this work, we created a control group that received no treatment or interruption whatsoever, and so, these were the least stressed. Mean sperm counts for test subjects was 49×10^6 in 72 hours compared to 99.5×10^6 in control rats. The mean percentage of sluggish and non-motile cells was 44% with acute treatment that further decreased to 35% in six weeks. However, there was evidence of improving testicular functions (the presence of immature spermatocytes and spermatids). There was improved sperm motility



Note the marked acute suppression of sperm cell count and motility

Figure 3: Sperm morphology, count, and motility after 72hrs. Note the marked acute suppression of sperm cell count and motility



There is suppression of cell count and cell motility but note that semen parameters are returning to normal (seen here as improving values).

Figure 4: Sperm morphology, count, and motility after 7 weeks from 56% acutely to 65.3% in the test rats after 6 weeks (65.3%) in comparison with 88% and 79% in the respective control rats. A decrease from 88% to 79% can be attributed to experimental stress. Sperm

morphologies were not affected by treatment but an assessment of the functional capacity of the cells can be confirmed (e.g. by mating treated rats with proven females). This may be more revealing but is outside the scope of this work.

The anti-spermatogenic effects of drugs and toxins may be due to their interference with Leydig cell function or steroidogenesis. Critical cellular processes that control steroidogenesis include the delivery of cholesterol into the mitochondria and steroidogenic enzyme expression in Leydig cells¹⁵. Disruption of Leydig cell functions, therefore, leads to reduced androgen production. *Ricinus communis*-linn contains agglutinin that binds to galactose, lactose, and their derivatives¹⁶. This reduces semen fructose, the major source of energy for spermatocytes, and may be responsible for the reduced motility (sluggishness) observed in this work. Nevertheless, this study revealed castor seed to be a potential male contraceptive agent that targets the process of spermatogenesis and motility of spermatocytes in particular, both in a reversible manner.

Conclusion

It is the right of the individual to access a preferred anti-fertility method. However, a method based on the male factor is simpler when compared to the diverse methods currently used by women. Moreover, the use of male contraceptives will reduce the stigma associated with the female on family planning. This work presents the possibility of a cheap, readily available, and potent, but a reversible male contractive agent of plant origin. The use of castor plant may be more acceptable culturally and religiously. It will not only improve female reproductive health but also alleviate the high female morbidity and mortality rates associated with conventional contraceptives.

Limitations

In future work, the effect of the extract on the ultrastructure of seminiferous tubules will be examined. Also, Leydig cell count and function will be evaluated. Moreover, the scope of this work does not address the effect of the extract on sperm capacitation.

Conflict of interest

The Authors have no conflict of interest in this work.

Acknowledgements

We thank Mr. John Ben, the Chief Technologist at the Anatomy Department, University of Jos, Nigeria for his technical assistance. We thank Prof C Isichei and the laboratory staff at the Faith Alive Hospital, Jos Nigeria.

References

- Chebet JJ, McMahon SA, Greenspan JA, Mosha IH, Callaghan-Koru JA, Killewo J, Baqui AH and Winch PJ. Every method seems to have its problems"- Perspectives on side effects of hormonal contraceptives in Morogoro Region, Tanzania. *BMC Women's Health* 2015; 15: 97-109. doi: 10.1186/s12905-015-0255-5.
- Kantorová V, Wheldon MC, Ueffing P and Dasgupta ANZ. Estimating progress towards meeting women's contraceptive needs in 185 countries: A Bayesian hierarchical modelling study. *PLoS Med.* 2020; 17(2): e1003026. Published online 2020 Feb 18. doi: 10.1371/journal.pmed.1003026.
- Amory JK. Male contraception. *Fertil Steril* 2016; 106(6): 1303-1309. doi:10.1016/j.fertnstert.2016.08.036.
- Datey SL, Gaur LN and Saxena BN. Vaginal bleeding patterns of women using different contraceptive methods (implants, injectable, IUDs, oral pills)—An Indian experience: An ICMR task force study. *Contraception* 1995; 51(3):155-165.
- WHO Department of Reproductive Health and Research. Unsafe abortion: global and regional estimates of the incidence of unsafe abortion and associated mortality in 2008. In: *Sexual and reproductive health 2011*; 6 Ed. ISBN: 978 92 4 150111 8.
- Grimes DA, Jones LB, Lopez LM and Schulz KF. Oral contraceptives for functional ovarian cysts. *Cochrane Database Syst Rev* 2009; CD006134.
- Odwe G, Obare F, Machiyama K and Cleland J. Which contraceptive side effects matter most? Evidence from current and past users of injectables and implants in Western Kenya. *Contracept X* 2020; 20(2): 100030. doi: 10.1016/j.conx.2020.100030. eCollection 2020.
- Cheng CY and Mruk DD. *Contraception* 2010; 82(5): 476–482. doi: 10.1016/j.contraception.2010.03.017. PMID: PMC4381878.
- United States Department of Agriculture (USDA). Plants Database. "*Ricinus Communis-Linn*". NRCS Plants Database. 2016. Team, Greensboro, NC 27401-4901 USA. <https://plants.sc.egov.usda.gov/home/>
- Heuzé V, Tran G, Hassoun P and Lebas F. Castor bean (*Ricinus Communis*) forage. Feedipedia, a programme by INRAE, CIRAD, AFZ and FAO 2020. <https://www.feedipedia.org/node/103>.
- Trease GE and Evans WC. Textbook of pharmacognosy. 12th Ed, Tindall and Co., London. 1983; 343-383.
- Irwin SD, Keen JN, Findlay JBC and Lord JM. The *Ricinus Communis* 2S albumin precursor: a single preproprotein may be processed into two different heterodimeric storage proteins. *Mol. Gen. Genet* 1990; 222: 400–408.
- Feng L, Li G, He Z, Han W, Sun J, Huang F, Di J and Chen Y. The ARF, GH3, and Aux/IAA gene families in castor bean (*Ricinus Communis-Linn*): Genome-wide identification and expression profiles in high-stalk and dwarf strains. *Industrial Crops and Products* 2019; 141: 111804.
- Hutchinson J and Dalziel J. M. Flora of West Tropical Africa. Crown Agents, (2nd Ed) (Revised by R.W.J. Keay) 1958. The Whitefriars Press, London and Tonbridge, Great Britain.
- Bradberry SM, Dickers KJ, Rice P, Griffiths GD and Vale JA. Review Ricin poisoning. *Toxicol Rev* 2003; 22(1): 65-70.
- Okwuasaba FK, Osunkwo MM, Ekwewchi, KE, Onwukeme K, Olayinka MO, Uguru VE and Das SC. Anticonceptive and oestrogenic effect of a seed extract of *Ricinus Communis-Linn* var. minor. *J. Ethnopharm* 1991; 34: 141–145.
- Okwuasaba FK, Das SC, Isichei CO, Ekwewchi MM, Onoruvwe O, Olayinka AO, Uguru VE, Dafur SJ, Ekwere EO and Parry O. Pharmacological Studies on the Antifertility Effects of Ricom-1013-J from *Ricinus Communis-Linn* var minor and Preliminary Clinical Studies on Women Volunteers. *Phytother Res* 1997; 11: 547-551.
- Osunkwo MM, Ibu J and Okwuasaba FK. Anticonceptive and oestrogenic effects of a medicinal plant (*Ricinus Communis-Linn*) in rats. In: 17th Annual Conference of the West African Society of Pharmacology, Port Harcourt Nigeria. 1988; 41–44.
- Usar JJ, Ekwere EO, McNeil RT and Okwuasaba KF. The Mechanism of Action and Effect on the Cervix of the Seed of *Ricinus Communis-Linn* Var. Minor (RICOM-1013-J) in Women Volunteers in Nigeria *J of Natural Sci Res.* 2016; 6(24): 64-70.
- Kabele-Toge, BB, Igwilo FT, Tayo F, Isamah GK and Eniojukan JF. Some toxicological and pharmacological studies on the extract of *Ricinus Communis-Linn* var. minor seeds with contraceptive properties. *Nig Cet Hosp Med* 1997; 7(4).
- McNeil RT, Okanlawon AO and Noronha CC. The Embryotoxic effect of a herbal extract with contraceptive properties. *J Med in the tropics* 2003; 5(1): 24-29.
- Makonnen E, Zerihun L, Assefa G and Rostom AA. Antifertility activity of *Ricinus communis* seed in female guinea pigs. *East Afr Med J.* 1999; 76(6): 335-337.

23. Farnsworth NR, Bingel AS, Cordell GA, Crane FA and Fong HH. Potential value of plants as sources of new antifertility agents. *J Pharm Sci* 1975; 64(4): 535-98.
24. Okwuasaba FK, Das SC, Isichei CO, Ekwenchi MM, Onoruvwe O, Olayinka AO, Uguru VE, Dafur SJ, Ekwere EO and Parry O. Pharmacological Studies on the Antifertility Effects of Ricom-1013-J from *Ricinus Communis-Linn* var minor and Preliminary Clinical Studies on Women Volunteers. *Phytother Res* 1997; 11: 547-551.
25. McNeil RT, Noronha CC, Kusemiju TO and Okanlawon AO. The Anti-ovulatory effect of a seed extract of *Ricinus Communis-Linn*. *Nig J Hlth Biomed Sci* 2003; 2: (1): 31-34.
26. Osunkwo MM, Ibu J and Okwuasaba FK. Anticonceptive and oestrogenic effects of a medicinal plant (*Ricinus Communis-Linn*) in rats. In: 17th Annual Conference of the West African Society of Pharmacology, Port Harcourt Nigeria 1988; 41-44.
27. Okwuasaba FK, Osunkwo MM, Ekwenchi KE, Onwukeme K, Olayinka MO, Uguru VE and Das SC. Anticonceptive and oestrogenic effect of a seed extract of *RICINUS COMMUNIS-LINN* var. minor. *J. Ethnopharm* 1991; 34: 141-145.
28. Isichei C, Das S, Ogunkeye O, Okwuasaba F, Uguru V, Onoruvwe O, Olayinka A, Dafur S, Ekwere E and Parry O. Preliminary clinical investigation of the contraceptive efficacy and chemical pathological effects of RICOM-1013-J of *Ricinus Communis-Linn* var Minor on women volunteers. *Phytotherapy Research* 2000; 14: 40-42.
29. McNeil RT, Noronha CC, Kusemiju TO and Okanlawon AO. The Anti-ovulatory effect of a seed extract of *RICINUS COMMUNIS-LINN*-linn. *Nig J of Health and Biomed Sci* 2003; 2: 31-34.
30. Ekwere OE, McNeil RT and Okwuasaba FK. The effect of *Ricinus Communis-Linn* (RICOM 1013-J) on semen parameters: A comparative study. *JPCS* 2011; 1: 8-11.
31. Okwuasaba FK, Osunkwo MM, Ekwenchi KE, Onwukeme K, Olayinka MO, Uguru VE and Das SC. Anticonceptive and oestrogenic effect of a seed extract of *Ricinus Communis-Linn* var. minor. *J. Ethnopharm* 1991; 34: 141-145.
32. Barrett KE, Barman SM and Boitano S. In: Brooks HL (Eds). *Ganong's Review of Medical Physiology*, 25th Ed. 2018. McGraw-Hill. <https://accessmedicine.mhmedical.com/content.aspx>.