

ORIGINAL RESEARCH ARTICLE

Socio-demographic and environmental determinants of child mortality in rural communities of Ogun State, Nigeria

DOI: 10.29063/ajrh2021/v25i5s.15

Dominic Azuh^{1,2,3}, Muiyiwa Oladosun^{1,2*}, Shalom Nwodo Chinedu^{3,4}, Akunna Ebere Azuh⁵, Emelda Duh¹, and Joy Nwosu¹

Department of Economics and Development Studies, Covenant University, Ota, Ogun State, Nigeria¹; Public-Private Partnership Research Cluster, Covenant University Centre for Research Innovation and Discovery (CUCRID), Ota, Ogun State, Nigeria²; Covenant University Public Health and Wellbeing Research Cluster (CUPHWERC)³; Department of Biochemistry, Covenant University, Ota, Ogun State, Nigeria⁴; Department of Business Management, Covenant University, Ota, Ogun State, Nigeria⁵

***For Correspondence:** Email: muiyiwa.oladosun@covenantuniversity.edu.ng

Abstract

Nigeria's under-five mortality was 132 per 1000 in 2018. The statistic makes Nigeria the country with the third-highest under-five mortality globally. It implies that the government may not achieve the Sustainable Development Goal (SDG) of 25 per 1000 births by 2030. This situation is of grave concern to policymakers and other stakeholders interested in the country's development. This study provides unique community micro-level information on child mortality determinants in rural communities where the country's health system is weakest. The study used a sample of 1350 pregnant women aged 20-44 who attended antenatal care in 22 health facilities in selected rural communities of Ogun State, South-west Nigeria. The multicollinearity diagnostics tests conducted between the dependent variable and predictors showed no abnormality in the values of the variance inflation factor, eigenvalues, and condition indexes. Logistics regression results showed that the socio-demographic characteristics such as the respondent's age, educational level, number of living children, and husband's education directly affected child mortality. In contrast, the husband has another wife had an indirect effect on child mortality. Environmental factors that directly impacted child mortality included the type of household toilet facility, source of water supply, and household waste disposal practices. These findings indicate that policies and programs to reduce child mortality in rural Nigeria must address socio-demographic and context-specific factors, especially at the community level. (*Afr J Reprod Health* 2021; 25[5s]: 158-169).

Keywords: Under-five mortality, socio-demographic, environmental factors, sustainable development goals

Résumé

La mortalité des moins de cinq ans au Nigéria était de 132 pour 1000 en 2018. Cette statistique fait du Nigéria le pays avec le troisième taux de mortalité des moins de cinq ans le plus élevé au monde. Cela implique que le gouvernement pourrait ne pas atteindre l'objectif de développement durable (ODD) de 25 pour 1000 naissances d'ici 2030. Cette situation est une grave préoccupation pour les décideurs politiques et les autres parties prenantes intéressées par le développement du pays. Cette étude fournit des informations uniques au niveau micro-communautaire sur les déterminants de la mortalité infantile dans les communautés rurales où le système de santé du pays est le plus faible. L'étude a utilisé un échantillon de 1350 femmes enceintes âgées de 20 à 44 ans qui ont assisté à des soins prénatals dans 22 établissements de santé dans certaines communautés rurales de l'État d'Ogun, au sud-ouest du Nigéria. Les tests de diagnostic de multicollinéarité effectués entre la variable dépendante et les prédicteurs n'ont montré aucune anomalie dans les valeurs du facteur d'inflation de la variance, des valeurs propres et des indices de condition. Les résultats de la régression logistique ont montré que les caractéristiques sociodémographiques telles que l'âge du répondant, son niveau d'instruction, le nombre d'enfants vivants et l'éducation du mari affectaient directement la mortalité infantile. En revanche, le mari a une autre femme a eu un effet indirect sur la mortalité infantile. Les facteurs environnementaux qui ont eu un impact direct sur la mortalité infantile comprenaient le type de toilettes domestiques, la source d'approvisionnement en eau et les pratiques d'élimination des déchets ménagers. Ces résultats indiquent que les politiques et programmes visant à réduire la mortalité infantile dans les zones rurales du Nigéria doivent tenir compte des facteurs sociodémographiques et spécifiques au contexte, en particulier au niveau communautaire. (*Afr J Reprod Health* 2021; 25[5s]: 158-169).

Mots-clés: Mortalité des moins de cinq ans, facteurs sociodémographiques, environnementaux, objectifs de développement durable

Introduction

The global under-five mortality rate fell to 39 deaths per 1,000 live births in 2018. Yet, children continue to face widespread regional disparities in their chances of survival, especially in sub-Saharan African countries¹. Childhood mortality has remained a significant challenge to public health in Nigeria and other developing countries. Under-five death is a fundamental measurement of a country's quality of life, health status, and development. In 2012, approximately half the world's estimated 6.6 million deaths in children aged less than five years occurred in sub-Saharan Africa, and Nigeria accounted for about 13% of these deaths². Previous studies have shown that half of the global under-five deaths occur in sub-Saharan Africa despite the region accounting for only one-fifth of the world's population of children^{3,4}.

Worldwide, sub-Saharan Africa has the highest under-five deaths globally, with 76 deaths per 1000, 14 times higher than the under-five mortality rate (U5MR) in developed countries^{5,6}. Nigeria's present under-five mortality is the eighth in the world⁷.

Notably, there has been a global decline in under-five mortality, mainly attributable to international initiatives and national programs that address the reduction of Under-5 mortality over the past years. By contrast, Nigeria is yet to maximize the positive effects of this global trend due to a lack of full comprehension of the factors affecting under-five mortality in the country. One of the United Nations' Sustainable Development Goals (SDGs) priority targets is to "end preventable deaths of newborns and under-five children by 2030"⁸. Nigeria's current under-five mortality rate of 132 deaths per 1000⁹ is high when compared to the expectation of the Sustainable Development Goal (SDG) of only 25 deaths per 1000 live births by 2030. Children under the age of five years are affected by socio-demographic and environmental factors prevalent in the middle- and low-income households in developing countries, including Nigeria.

According to Emenike *et al.*¹⁰, potable water is essential to maintain a clean environment and ensure good health. Conversely, limited access

to safe drinking water supply, quality sanitation facilities, and unhealthy hygiene practices promote the spread of waterborne diseases, which causes 6.3% of the deaths recorded around the world¹¹. Young children are the first to get sick and die from waterborne and sanitation-related illnesses, including diarrheal diseases and malaria². The UNICEF also observed that children in sub-Saharan Africa (SSA) are 14 times more likely to die before the age of five than children from developed regions¹². Poor environmental sanitation conditions contribute significantly to the high prevalence of infectious diseases, including malaria, cholera, typhoid, diarrhea, acute respiratory infection, and tuberculosis, among others¹³. By 2011, an estimated one billion people still used open defecation, and 185 million people relied on surface water for drinking¹².

In a study in Nigeria, Olukanni *et al.*¹⁴ postulated that inadequate waste management and poor sanitation practices had become a significant concern in Nigeria and many developing countries in SSA. According to WHO/UNICEF¹⁵, an estimated 9% of people worldwide lack access to potable water, and 2.4 billion cannot adequately access quality sanitation facilities despite the coordinated global efforts to actualize the SDG targets. Research evidence indicates that diarrheal illnesses in children account for the high rate of under-five mortality in Nigeria¹⁶. The negative consequences of household and environmental factors on under-five children's morbidity and mortality are documented well¹⁷.

There is substantive evidence indicating that household environmental conditions such as drinking water sources, types of primary floor materials, and ambient air quality are determinants of childhood mortality^{9,18,19}. Studies have also shown that mothers with formal education have reduced under-five mortality compared to those with no formal education^{7,20,21}. Household environmental characteristics have a significant impact on child mortality as lower mortality rates are experienced in households with access to immunization, proper refuse and solid waste disposal facilities, and access to drinking water and sanitation facilities⁴. Potable water is essential to maintain a clean environment, ensure good health,

and a healthy nation at¹⁰. Children are the first to get sick and die from waterborne and sanitation-related illnesses include diarrheal and malaria diseases².

Providing safe drinking water and access to improved sanitation within the household environment can reduce the risk of morbidity and mortality among under-five aged children²². According to UNICEF and WHO reports, insufficient water, sanitation, and hygiene issues account for many illnesses. The most common waterborne disease like diarrhea, having an annual incidence of 4.6 billion episodes and cause 2.2 million deaths every year in developing countries². As in many developing countries like Nigeria, socio-demographic and environmental factors are crucial to understanding the states' persistently relatively high child mortality. Literature suggests that under-five death is inversely related to mother's age; that is, as mother advances in age, child mortality rate reduces^{23,24}. Mothers who attained higher educational levels had lower under-five mortality than their counterparts who had no education²⁴ and Under-five mortality is higher in families with limited access to drinking water.

Children born in households with inadequate toilet facilities experience the highest mortality rate (41.0%) than those born in homes with improved bathrooms (30.4%)²⁵. Furthermore, Mutunga²⁶ reported that among the ten identified leading mortality risks in developing countries, unsafe water, sanitation, hygiene, and smoke from solid fuels were among the most threatening. Mesike and Mojekwu⁴ reported that about 3% of the resulting deaths are attributable to environmental risk factors, and child deaths account for about 90% of the total deaths.

Child mortality reduction has become a common concern of all government tiers in Nigeria, non-governmental organizations, and international agencies. It is imperative to accelerate progress in preventing child deaths as current trends predict that close to 52 million under-five aged children will die between 2019 and 2030¹. Understanding the socio-demographic and environmental factors influencing under-five mortality is significant in providing useful insights that would help formulate effective public health interventions, which can scale down the high mortality among under-five children leading to the achievement of SDG 3. It is

against this background that this community-based study focused on the Ifo local government area of Ogun State. It examined socio-demographic and environmental factors influencing under-five child mortality at the community level and generating appropriate policy and programmatic actions.

Methods

The paper was on a cross-sectional hospital-based survey data collected using a multi-stage sampling technique to select the respondents. Ifo Local Government Area (LGA) of Ogun State, Nigeria, was the study's location. The first stage of sampling involved a purposive selection of one LGA from 20 in the state. An LGA with a reasonably large population, high child mortality, and proximity to the research team's location were purposive selection criteria. In the second sampling stage, a total of 29 health facilities were in the LGA, of which 22 facilities were selected systematically with a random start and sampling fraction of 2, amounting to 75.9% of the total. This study's analysis unit was pregnant women who attended antenatal care and gave birth to at least one child in the last three years preceding the survey. Guided by the patients' list in the health facilities used as the sampling frame, pregnant women were clustered according to their clinic days. The team interviewed all women present during the clinic days that fall within the study period were interviewed.

The study team conducted fieldwork between May 1 and July 31, 2018. Overall, 1350 pregnant women respondents (6.76%) constituted the sample from a base population of 19,964 pregnant women who registered at the 29 health facilities at the time of the study. The research instrument was validated by two experts, a demographer, and health care personnel. In addition to the Cronbach Alpha index, which yielded 0.75, the data collection instrument was pretested, and questions were modified to perfect its reliability. The study adopted part of the 2013 Nigeria Demographic and Health Survey (NDHS) questionnaire to suit the local context. The structured questionnaire captured information on demographic characteristics, child mortality, and survival characteristics. The survey questionnaire was conducted by a team of experienced nurses and

matrons trained specially for the study before its commencement. Besides, they were fluent in English, Yoruba, and the local dialect of the community members.

Variables operationalization

The study conceptualized that socio-demographic and environmental factors have direct and indirect relationships with child mortality. The study data obtained went through transformation and recoding using SPSS Version 20, and categorization and recoding were based mostly on available literature (e.g., NDHS categories) and the decision to reduce empty cells thus, strengthening degrees of freedom, analysis, and the results. Child mortality is the dependent variable in this study and was captured in the survey as ever lost any children aged 1-5 years and responses categorized as 1 = yes, and two = no. In this study, socio-demographic predictors of child mortality were the age of respondent, age at delivery, religion, marital status, had co-wife, education, occupation, and the number of living children. Other socio-demographic characteristics included in the study were the spouse's education and occupation.

Age of respondent was categorized as 20-24 = 1, 25-30 = 2, 31-40 = 3, 30 and older = 4. Age that respondent had their first child was obtained in single years, and regrouped as 20 or less = 1, 21-24 = 2, 25-29 = 3, and 30 or older = 4. Religion was coded as traditional = 1, Islam = 2, and Christianity = 3, marital status recategorized into single/divorced/separated/widowed = 1, and married = 2, and co-wife, captured as whether husband had another wife was coded as yes = 1, no = 2. Education another socio-demographic factor was recoded as none/primary = 1, secondary = 2, post-secondary/professional = 3; occupation was captured as not working = 1, self-employed = 2, civil servant = 3, and private sector employee = 4; children ever born was a continuous variable grouped as 1 or 2 = 1, 3 or 4 = 2, and 5 or more = 3; while number of living children which captures current fertility at time of survey was regrouped in similar fashion as children ever born. Other socio-demographic variables included in the study were spouse's education recorded as none/primary = 1, secondary = 2, and post-secondary/professional = 3; and spouse's occupation recorded as not working

= 1, self-employed = 2, civil servant = 3, and private sector employee = 4.

Environmental predictors of child mortality are conceptualized in this study as those that affect living conditions with consequential effects on child health status. Key environmental factors considered in this study were the type of house lived in, the number of persons in a household, the type of toilet in the household, the main source of water supply, and the household waste disposal system⁴. The type of household respondent lived in was categorized as mud/grass/hut = 1, one room = 2, room and parlor = 3, two/three bedroom flat = 4, and detached house/mansion = 5. Number of persons in household was captured in single years and then categorized as 1 or 2 = 1, 3-4 = 2, 5-6 = 3, and 7 and above = 4. Type of toilet in household was measured as field/bush = 1, bucket toilet = 2, pit latrine = 3, and flush toilet = 4. Main source of household water supply was measured as river/steam = 1, covered well = 2, tanker/truck = 3, borehole = 4, and tap = 5. Moreover, the household waste disposal system was measured as government collection = 1, private agency = 2, disposal within the compound (including burying and burning) = 3, and unauthorized dumpsite = 4.

Results

Sample statistics

Table 1 shows that most of the women interviewed were aged 30 or younger (61.4%), and most had their first child before reaching age 30. They were mainly Christians (59.1%), married (84.5%), and their husbands had another wife (76.2%). The majority of the respondent had secondary or higher education (85%), were self-employed (72.8%), and about half (51.7%) had three or more children alive. Most women's husbands had at least a secondary education (88.8%) and were self-employed (75.2%). On environmental factors, most study respondents and their household (68.5%) lived in a room and parlor/lower-level accommodation and were at least three persons in such housing unit (81.9%). Respondents reported that their toilets were mostly flushed (67.6%), and their primary source of water supply was borehole/tap (81.9%). Most health facilities were at trekkable distance to the respondents (75.3%). And most households of

Table 1: Percentage distribution of pregnant women who lost children by socio-demographic and environmental factors

Variables	Frequency	%
Total N	1350	
Age of respondent		
20-24	283	21.0
25-30	546	40.4
31-40	391	29.0
41 and above	130	9.6
Age of respondent at delivery		
20 or less	159	12.1
21-24	473	35.9
25-29	485	36.8
30 and above	202	15.3
Religion of respondent		
Traditional	87	6.5
Islam	463	34.4
Christianity	794	59.1
Marital status		
single-divorced-separated-widowed	209	15.5
Married	1141	84.5
Husband had another wife		
Yes	321	23.8
No	1029	76.2
Education of respondent		
none-primary	203	15.0
Secondary	536	39.7
post-sec-professional	611	45.3
Respondent occupation		
not working	147	14.2
self-employed	751	72.8
civil-servant	134	13.0
Number of living children		
1 or 2	652	48.3
3 or 4	489	36.2
5 or more	209	15.5
Spouse education		
none-primary	152	11.3
Secondary	541	40.1
post-sec-professional	657	48.7
Spouse occupation		
not working	86	8.3
self-employed	781	75.2
civil-servant	171	16.5
Type of house living in		
mud-grass-hut	100	7.4
one room	379	28.1
room and parlor	445	33.0
two-three bedroom flat	369	27.3
detached house-mansion	57	4.2
Number of persons living in a house		
2 or 1	243	18.0
3-4	689	51.0
5-6	353	26.1
7 and above	65	4.8
Type of toilet facility for household		
field-bush/bucket toilet	137	10.2
pit-latrine	300	22.3

Variables	Frequency	%
flush toilet	910	67.6
Main source of water supply		
river-stream	132	9.8
covered well	28	2.1
tanker-truck	84	6.2
Borehole	497	36.8
Tap	608	45.1
Distance of house to health facility		
not far/trekkable	622	46.1
far but trekkable	394	29.2
very far/don't know	334	24.7
Household waste disposal practices		
govt collection	679	50.3
private agency	326	24.1
disposal within comp (burying or burning)	236	17.5
unauthorized dumpsite	109	8.1
Ever lost any child(ren) aged 1-5 years		
No	1055	78.2
Yes	294	21.8
Total N	1350	

respondents disposed of their waste through government/private arrangement (74.4%). About one-fifth (21.8%) of the respondents reported the death of children aged 1-5 years.

Bivariate results

Table 2 shows the association between the dependent variable lost children aged 1-5 by socio-demographic and environmental factors. Chi-Square statistical technique was used to test the association between the dependent and independent variables at .1, .05, .01, and .001 levels. In Table 2 socio-demographic factors that had significant association with child mortality were age of respondent at delivery (p-value = .099), religion of respondent (p-value = .000), marital status (p-value = .001), husband had another wife (p-value = .000), education of respondent (p-value = .009), and number of living children (p-value = .000). Environmental factors associated with the death of children aged 1-5 were; the type of house lived in (p-value = .054), and household medium of waste disposal (p-value = .001).

Multivariate results

A crucial step in the multivariate level analysis is collinearity diagnostics. Appendix 1 shows the test results on collinearity between socio-demographic

Table 2: Percentage frequency distribution of pregnant women who lost any children aged 1-5 by socio-demographic and environmental factors

Ever lost any child(ren) aged 1 – 5 years			
Variables	No (%)	Yes (%)	p-value
Age of respondent			
20-24	22.0	17.0	
25-30	40.7	39.8	
31-40	28.3	31.3	
41 and above	9.0	11.9	.140
Age of respondent at delivery			
20 or less	12.9	9.0	
21-24	35.8	35.8	
25-29	37.0	36.1	
30 and above	14.3	19.1	.099
Religion of respondent			
Traditional	5.0	11.6	
Islam	34.8	32.9	
Christianity	60.1	55.5	.000
Marital status			
single-divorced-separated-widowed	13.7	21.4	
Married	86.3	78.6	.001
Husband had another wife			
Yes	21.1	33.0	
No	78.9	67.0	.000
Education of respondent			
none-primary	14.6	16.7	
Secondary	41.9	32.0	
post sec-professional	43.5	51.4	.009
Respondent occupation			
not working	14.8	12.0	
self-employed	72.3	74.5	
civil-servant	12.9	13.4	.576
Number of living children			
1 or 2	50.9	39.1	
3 or 4	37.3	32.7	
5 or more	11.8	28.2	.000
Spouse education			
none-primary	11.4	10.9	
Secondary	39.6	41.5	
post sec-professional	49.0	47.6	.843
Spouse occupation			
not working	8.5	7.4	
self-employed	74.9	76.4	
civil-servant	16.6	16.2	.842
Type of house living in			
mud-grass-hut	6.4	11.2	
one room	28.2	27.9	
room and parlor	34.1	28.9	
two-three bedroom flat	27.2	27.6	
detached house-mansion	4.2	4.4	.054
Number of persons living in a house			
2 or 1	18.8	15.3	
3-4	51.6	49.3	
5-6	24.9	30.6	
7 and above	4.7	4.8	.204
Type of toilet facility for household			
field-bush/bucket toilet	10.3	9.6	
pit-latrine	21.9	23.5	
flush toilet	67.8	66.9	.816

Ever lost any child(ren) aged 1 – 5 years			
Variables	No (%)	Yes (%)	p-value
Main source of water supply			
river-stream	10.0	9.2	
covered well	1.7	3.4	
tanker-truck	5.7	7.8	
Borehole	37.1	36.1	
Tap	45.5	43.5	.259
Distance of house to health facility			
not far/trek able	46.8	43.2	
far but trek able	29.0	29.9	
very far/don't know	24.2	26.9	.496
Household waste disposal practices			
govt collection	49.5	53.4	
private agency	25.3	20.1	
disposal within comp (burying or burning)	18.5	13.6	
unauthorized dumpsite	6.7	12.9	.001
Total N		1350	

Table 3: The odds that pregnant women lost any children by socio-demographic and environmental determinants

Variable	Model 1 (only socio-demographic factors)				Model 3 (All predictors)			
	Exp(B)	Lower	Upper	Sig.	Exp(B)	Lower	Upper	Sig.
Socio-demographic factors								
Age of respondent								
20-24	1.00				1.00			
25-30	1.454	.874	2.419	.149	1.421	.840	2.406	.190
31-40	1.874	1.083	3.244	.025	1.799	1.016	3.185	.044
41 and above	.825	.375	1.812	.632	.796	.354	1.792	.582
Age of respondent at delivery								
20 or less	1.00				1.00			
21-24	1.385	.729	2.633	.320	1.304	.676	2.515	.429
25-29	1.135	.580	2.222	.712	1.102	.554	2.194	.782
30 and above	1.296	.595	2.822	.514	1.233	.551	2.760	.610
Religion of respondent								
Traditional	1.00				1.00			
Islam	.608	.319	1.158	.130	.525	.268	1.029	.061
Christianity	.702	.369	1.336	.281	.582	.296	1.141	.115
Marital status								
single-divorced-separated-widowed	1.00				1.00			
Married	.754	.473	1.203	.237	.796	.487	1.299	.360
Husband had another wife								
Yes	1.00				1.00			
No	.611	.400	.934	.023	.577	.370	.901	.016
Education of respondent								
none-primary	1.00				1.00			
Secondary	.692	.408	1.173	.171	.658	.379	1.141	.136
post sec-professional	1.360	.749	2.468	.312	1.367	.735	2.541	.324
Respondent occupation								
not working	1.00				1.00			
self-employed	1.009	.578	1.760	.975	.920	.518	1.637	.778
civil-servant	1.031	.484	2.195	.938	.895	.408	1.967	.783
Number of living children								
1 or 2	1.00				1.00			
3 or 4	.986	.653	1.488	.946	.906	.585	1.403	.659
5 or more	3.095	1.979	4.841	.000	2.875	1.748	4.727	.000
Spouse education								
none-primary	1.00				1.00			
Secondary	1.694	.929	3.088	.085	1.743	.936	3.246	.080
post sec-professional	1.464	.747	2.872	.267	1.436	.711	2.900	.313

Variable	Model 1 (only socio-demographic factors)				Model 3 (All predictors)			
	Exp(B)	Lower	Upper	Sig.	Exp(B)	Lower	Upper	Sig.
Spouse occupation								
not working	1.00				1.00			
self-employed	1.314	.652	2.648	.445	1.311	.631	2.727	.468
civil-servant	1.006	.441	2.293	.989	1.059	.447	2.511	.897
	Model 2							
	Exp(B)	Lower	Upper	Sig.				
Type of house living in								
mud-grass-hut	1.00				1.00			
one room	.617	.373	1.020	.060	.817	.419	1.593	.553
room and palour	.501	.304	.828	.007	.830	.416	1.655	.597
two-three bedroom flat	.568	.338	.955	.033	.778	.369	1.643	.511
detached house-mansion	.592	.272	1.292	.188	.660	.205	2.124	.486
Number of persons living in the house								
2 or 1	1.00				1.00			
3-4	1.256	.856	1.844	.244	1.107	.664	1.846	.696
5-6	1.736	1.139	2.646	.010	1.733	.970	3.095	.063
7 and above	1.226	.610	2.463	.567	.648	.262	1.598	.346
Type of toilet facility for household								
field-bush/bucket toilet	1.00				1.00			
pit-latrine	1.474	.868	2.503	.151	2.104	1.037	4.271	.039
flush toilet	1.400	.848	2.312	.188	2.506	1.269	4.950	.008
Main source of water supply								
river-stream	1.00				1.00			
covered well	2.144	.862	5.330	.101	1.545	.539	4.428	.418
tanker-truck	1.454	.734	2.880	.283	1.233	.520	2.923	.635
Borehole	1.033	.621	1.718	.901	.773	.419	1.429	.412
Tap	1.008	.618	1.643	.975	.752	.419	1.350	.340
Distance of house to health facility								
not far/trekkable	1.00				1.00			
far but trekkable	1.182	.862	1.620	.300	1.122	.730	1.727	.599
very far/don't know	1.132	.810	1.581	.467	1.111	.714	1.730	.640
Household waste disposal practices								
govt collection	1.00				1.00			
private agency	.754	.537	1.059	.104	.819	.525	1.278	.379
disposal within comp (burying or burning)	.615	.411	.920	.018	.652	.383	1.109	.114
unauthorised dumpsite	1.843	1.171	2.900	.008	1.671	.902	3.095	.103

Note: Level of Significance; . *p ≤ .05, **p ≤ .01, ***p ≤ .001.
Model 1: Chi-square = 61.192, -2 Log likelihood = 834.359^a, Nagelkerke R Square = .106
Model 2: Chi-square = 38.560, -2 Log likelihood = 1371.445^a, Nagelkerke R Square = .044.
Model 3: Chi-square = 83.665, -2 Log likelihood = 810.942^a, Nagelkerke R Square = .143

and environmental factors. Except in a few cases, i.e., six of 256 variance proportion values, the majority were less than .5, and results of statistical test significant. None of the six variance proportion values were corroborated by results of other collinearity indicators, i.e., VIF, CI, Eigenvalues, and tolerance.

Table 3 (Model 1) examined relationships between the dependent variable, lost any children aged 1-5, and socio-demographic predictors. Model

2 presents the relationships between the dependent variable and only environmental factors. Model 3 shows the relationships between the dependent variable and socio-demographic and environmental factors both included in the equation. Thus, the three blocks of models enabled teasing out direct and indirect effects of socio-demographic and environmental predictors of child mortality. The strength of the relationships regarding point estimates was tested at four levels-- .1, .05, .01, and

.001. Also, confidence intervals in Table 3 depict the range of acceptance of the estimated population parameters at 95% confidence. The summary results for the three models are presented in the last panel of Table 3. The results showed that Model 3 compared with Models 2 and 1 is best fitted with a higher Chi-square value (83.67, vs. 38.56 and 61.19, respectively), and lowest -2 log-likelihood (810.94 vs. 1371.44 and 834.36, respectively), and a higher proportion of explained variation (14.3% vs. 11%, and 4.4%, respectively).

This section's findings focussed mainly on significant outcomes in Model 3 of Table 3, which is the best fit and has all explanatory variables. Model 3 shows that child mortality had indirect relationships with the type of house lived in and household waste disposal practices. Moreover, child mortality directly links with respondent age, religion; the husband had another wife, number of living children, husband's education, number of persons who lived in the house, and type of toilet facility for respondents' household.

Relationships between child mortality and socio-demographic factors

The odds of child mortality were about two times higher for respondents aged 31-40 compared to their younger counterparts aged 20-24 (OR = 1.79, [CI = 1.02, 3.18]). The odds of losing children was significantly lower for Muslim respondents compared to their traditionalist counterparts (OR = .525, [CI = .268, 1.03]), and it was lower for those whose husband had another wife compared to their traditionalist counterpart (OR = .577, [CI = .370, .901]). As expected, the odds of child mortality were higher for respondents who had five or more living children than those with only one or two living children (OR = 2.87, [CI = 1.75, 4.73]). The odds of child mortality were higher for women whose husbands had secondary education compared to those whose husbands had none or primary (OR = 1.74, [CI = .936, 3.25]).

Relationships between child mortality and environmental determinants

Concerning environmental factors, findings showed that the odds of child mortality were about twice for respondents who reported having five or

six persons in their household than their counterparts who reported two or one (OR = 1.73, [CI = .970, 3.09]). The relationship between child mortality and type of toilet facility in the household was contrary to expectation. The odds of child mortality for respondents with pit-latrine was twice that of their counterparts who used field-bush/bucket as toilet facility (OR = 2.10, [CI = 1.04, 4.27]). Also, the odds of a child aged 1-5 dying for respondents who used a flush toilet in their household was 2.5 times compared to their colleagues who used field-bush/bucket as a waste facility (OR = 2.51 [CI = 1.27, 4.95]).

Discussion

This study examined socio-demographic and environmental factors influencing child mortality in a rural context in Nigeria, where evidence of this nature is rare and virtually non-existent. It conducted a hospital-based study collecting health and related information from 1350 pregnant women attending an ante-natal clinic at the survey time. Findings provide insights into direct and indirect socio-demographic and environmental factors affecting child mortality among the communities studied. The importance of these results for programming is prioritizing intervention for what should come first by order of importance and perhaps impact. Factors that directly affected child mortality were the respondent's age, religion, family type (monogamous vs. polygamous), number of living children, and husband's education. Other independent variables that directly affect child mortality are the number of persons in a house and the type of household toilet facility. These predictors of child mortality should be considered before other factors like the type of home lived in and household disposal practices that have indirect effects on child mortality. The distinction between direct and indirect predictors of child mortality is a unique contribution of this paper. The findings that directly affect child mortality may likely be more efficient in reducing child mortality than the indirect effects. Thus, policy and program implementation should be prioritized, considering factors with direct effects before those with indirect effects. Age of respondents, religion, no of co-wives, the number

of living children, spouse's education are socio-demographic factors that should be considered in a future intervention with the objective of child mortality reduction in the locality and perhaps other rural communities in the country. Policy and program intervention need to target older women, especially those who have many children, with information that will encourage them to take good care of their children, especially those aged between one and five. It will be necessary to customize program intervention based on region and focus more on traditional women to change their behavior on better care for their children. It was entirely unexpected that spouses' education had adverse effects on child mortality, and women's education was not significant as in other studies^{7,20}. The results may be due to male hegemony and influence in a typical Nigerian household, especially in the rural areas. Thus, the decisions of educated men in community settings are perhaps, more respected than those of their counterparts who were not educated. Further research will be necessary to shed more insights into this unexpected negative outcome of this study.

A critical environmental predictor of child mortality in this study is the number of persons living in a household. The number of persons in a house is an indicator of density or crowded home, which implies more pressure on available resources and facilities such as potable water²⁶ and increased risk factors including poor hygiene and unclean environment^{4,10}. In this circumstance, children can easily be at the receiving end of the ladder, resulting in increased mortality risk²⁵. Policy and programs should address this issue by educating families on the danger of overcrowded households and the possible consequences. Also, government decision-makers at the state level and other interested stakeholders should increase access to affordable housing, potable water, and other amenities at the community level, thus, improving quality of life and reduce child mortality in the long-run.

Another interesting finding of this study is that women who reported that their home has flush toilets had higher odds of child mortality than their counterparts who defecated in the field/bush or used bucket toilets. While this paper may not provide specifics on the reasons for this result, it is appropriate to note that the result is suggestive that

while having a flushed toilet may be necessary for a household, it is not sufficient to attain good hygiene. Anecdotal evidence suggests that households in rural communities of this study are likely to have more flushed toilet facilities without adequate water supply than with water supply. Therefore, for the dividends of using the flushed toilet to be realized in households in such rural settings, it will be necessary to complement a constant and adequate supply of water to ensure that flushed toilets are clean at all-time^{9,10,18,19}. Otherwise, the flushed toilets may serve as sources of diarrhea, one of the killer diseases of children¹⁰. One limitation of this study is that the water supply source's information was not enriched by constant water supply or household utilization information. Studies in the literature have examined unsafe water and its negative impact on child mortality^{10,26-32}; what is yet to be teased out or to receive significant attention is the relationships between the flushed toilet, water supply, and child mortality. Policy and program intervention geared to achieving SDG 3.2 by 2030 must ensure adequate supply and access to portable water supply^{33,34} in Ogun State and Nigeria's rural communities.

Ethical consideration

The research team got approval from the Covenant University management to conduct the study. The team also applied to the Ifo Local Government Area (LGA) Chairman and was given tacit approval to conduct the study in the selected health facilities in the LGA. The study did not involve any activity that may cause harm or risk to human life. However, the study team applied standard research ethics, including informed consent, willingness to participate, and anonymity of the respondent.

Conclusion

This study examined vital socio-demographic and environmental factors affecting child mortality in selected rural communities in Nigeria. The main factors to improve child mortality in the household are direct and indirect, based on their predictive paths. The indirect factors are the type of house lived-in and household waste disposal practices. The direct predictors of child mortality for intervention purposes are the age of women,

religion, kind of marriage, number of living children, husband's education, and type of household toilet facility. These direct factors should be given priority in policies and programming focused on improving child mortality in rural areas in Nigeria. Prioritizing programs based on the direct and indirect factors influencing child mortality will go a long way to achieving SDG 3.2, which focuses on ending newborn and child mortality in the country by 2030.

Acknowledgement

The authors are grateful for publication funding support made available by Covenant University through the Covenant University Centre for Research, Innovation and Discovery (CUCRID).

References

1. Suggested citation: United Nations Inter-agency Group for Child Mortality Estimation (UN IGME), 'Levels & Trends in Child Mortality: Report 2019, Estimates developed by the United Nations Inter-agency Group for Child Mortality Estimation', United Nations Children's Fund, New York, 2019.
2. UNICEF (2012). WHO Countdown to 2015 Decade Report (2000-2010): Taking Stock of Maternal, Newborn, and Child Survival.
3. Smith EG. Maternal Schooling and Child Mortality in Nigeria: The importance of the Actual curriculum [Online] Available. 2010: Princeton.edu/download.aspx?submissionId=100377.
4. Mesike CG and Mojekwu JN. Environmental determinants of child mortality in Nigeria. *Journal of Sustainable Development*. 2012; 5(1): 65-75.
5. Adebawale AS, Adeniyi FF, Oyewale M, Taiwo O, Rotimi FA and Martin EP. Parental Educational homogamy and under-five mortality in sub-Saharan Africa: Clarifying the association's intricacy. *Scientific African*. 2020; 7 e00255, 1-17
6. United Nations Inter-Agency Group for Child Mortality Estimation (UN IGME). 2018. 'Levels & Trends in Child Mortality: Report 2018, Estimates developed by the United Nations Inter-Agency Group for Child Mortality Estimation. United Nations International Children's Emergency Fund (UNICEF): New York.
7. Yaya S, Bishwajit G, Okonofua F and Uthman OA. Under-five mortality patterns and associated maternal risk factors in sub-Saharan Africa: A multi-country analysis. *PLoS ONE*. 2018; 13(10): e0205977. Pp. 1-14. <https://doi.org/10.1371/journal.pone.0205977>.
8. United Nations. 2015. Transforming Our World: The 2030 Agenda for Sustainable Development. A/RES/70/1. www.sustainabledevelopment.un.org.
9. National Population Commission (NPC) and ICF. 2019. Nigeria Demographic and Health Survey, 2018. Key Indicator Report. Abuja and Rockville, Maryland, USA: NPC & ICF.
10. Emenike CP, Tenebe IT, Omole DO, Ngene BU, Oniemayin BI, Maxwell O and Onoka BI. Accessing Safe Drinking Water in sub-Saharan Africa: Issues and Challenges in South-West, Nigeria. *Sustainable Cities and Society*. 2017; <http://dx.doi.org/10.1016/j.scs.2017.01.005>.
11. Omole DO and Ndambuki JM. Sustainable Living in Africa: Case of Water, sanitation, Air Pollution, and Energy. *Sustainability*. 2014; 6(8), 5187-5202.
12. WHO, UNICEF and Joint Water Supply & Sanitation Monitoring Program (2013). Progress on Sanitation and Drinking Water: 2013 Update. Geneva, Switzerland: WHO and UNICEF.
13. Cleveland M, Kalaness M and Laroche M. It is not easy being Green. Exploring Green Creeds, Green Deeds, and Internal Environmental Locus of Control. *Psychology and Marketing*. 2012 29(5), 293-305.
14. Olukanni DO, Azuh DE, George TO, Ajayi MP and Emenike PC. The Relevance of Policy and Practice on Sanitation Effort in Developing Nations: The Experience of a Semi-urban city in southwest Nigeria. 2014; Proceedings of ICERI 2014 Conference.
15. WHO and UNICEF (2015). 25 Years Progress on Drinking Water and Sanitation 2015 Update and MDG Assessment. [Assessed 16/05/2016 at http://www.wssinfo.org/fileadmin/user_upload/resources/JMP-Update-2015_English.pdf]
16. Adewemimo A, Kalter HD, Perin J, Koffi AK, Quinley J and Black RE. Direct Estimates of Cause-specific Mortality Fractions and Rates of Under-five Deaths in the Northern and Southern Regions of Nigeria by Verbal Autopsy Interview. *PLoS ONE*. 2017; 12, e0178129.
17. Brown J, Sandy C and Jeroen HJE. Water, Sanitation, Hygiene, and Enteric Infections in Children. *Global Child Health, Arch Dis Child*. 2013; 98: 629-634.
18. Adebawale SA, Morakinyo OM and Ana GR. Housing materials as predictors of under-five mortality in Nigeria: evidence from 2013 demographic and health survey. *BioMed Central (BMC) pediatrics*. 2017; 17(1), 30.
19. Lelieveld J, Haines A and Pozzer A. Age-dependent health risk from ambient air pollution: a modeling and data analysis of childhood mortality in middle-income and low-income countries. *The Lancet Planetary health*. 2018; 2(7), e292-e300.
20. Fasina F, Oni G, Azuh D and Oduaran A. Impact of Mother's Socio-demographic Factors and Antenatal Clinic Attendance on Neonatal Mortality in Nigeria. *Cogent Social Sciences*. 2020; 6(1), 1747328.
21. Adepoju AO, Akanni O and Falusi AO. Determinants of child mortality in Rural Nigeria. *World Rural*

- Observations. 2012; 4(2), 38-45.
22. WHO (2009). World Health Statistics, 2009. Geneva: The World Health Organization.
 23. Gebretsadik S and Gabreyohannes E. Determinants of Under-Five Mortality in High Mortality Regions of Ethiopia: An analysis of the 2011 Ethiopia Demographic and Health Survey Data. *International Journal of Population Research*. 2016; <http://dx.doi.org/10.1155/2016/1602761>.
 24. Ezech OK, Agho KE, Dibley MJ, Hal JJ and Page AN. Risk factors for post neonatal, infant, child, and under-5 mortality in Nigeria: a pooled cross-sectional analysis. *BMJ Author Services*. 2019; Vol. 5, Issue 3. <http://bmjopen.bmj.com/content/5/3/e006779>.
 25. Adeolu MO, Akpa OM, Adeolu AT and Aladeniyi IO. "Environmental and Socioeconomic Determinants of Child Mortality: Evidence from the 2013 Nigerian Demographic Health Survey." *American Journal of Public Health Research*. 2016; vol. 4, no. 4: 134-141. doi: 10.12691/ajphr-4-4-3.
 26. Mutunga CJ. Environmental determinants of child mortality in Kenya. UNU-WIDER Research Paper No 2007/83. Helsinki: United Nations University World Institute for Development Economics Research. 2007;
 27. Yaya S, Hudani A, Udenigwe O, Shah V, Ekholuenetale M and Bishwajit G. Improving Water, Sanitation and Hygiene Practices, and Housing Quality to Prevent Diarrhea among Under-Five Children in Nigeria. *Trop. Med. Infect. Dis*. 2018; 3, 41; pp. 1-11. doi: 10.3390/tropicalmed302004
 28. World Health Organization. 2008. World Health Organization and the United Nations children's fund Joint Monitoring Programme for water supply and sanitation. Progress on drinking water and sanitation: special focus on sanitation. In the World Health Organization and the United Nations children's fund Joint Monitoring Programme for water supply and sanitation (JMP). Progress on drinking water and sanitation: special focus on sanitation. World Health Organization.
 29. WHO (2013). Health Topics: Sanitation. <http://www.who.int/topics/sanitation/en/>
 30. WHO (2011). Ten Facts on Sanitation. <http://www.who.int/features/factfiles/sanitation/en/index.html>
 31. WHO, UNICEF, and Joint Water Supply & Sanitation Monitoring Program (2013). Progress on Sanitation and Drinking Water: 2013 Update. Geneva, Switzerland: WHO and UNICEF.
 32. WHO and UNICEF (2012). Progress on Drinking Water and Sanitation 2012 Update in Joint Monitoring Programme for Water Supply and Sanitation.
 33. United Nations. 2015. Transforming our world: The 2030 agenda for sustainable development. Retrieved March 31, 2017, from https://www.un.org/.../120815_outcome-document-of-Summit-for-adoption-of-the-p
 34. United Nations General Assembly. Transforming our world: the 2030 Agenda for Sustainable Development. New York: United Nations, 2015. http://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E (accessed January 16, 2018).

Appendix

Appendix I: Collinearity Diagnostics on number of children ever lost aged 1-5 by socio-demographic and environmental factors

Variance Proportions																Sig	Tol.	VIF	Eigen value	Cond index
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16					
1.	.03	.00	.00	.00	.01	.02	.00	.08	.01	.00	.01	.02	.01	.02	.27	.962	.793	1.261	.261	7.709
2.	.09	.04	.00	.00	.00	.00	.00	.16	.00	.00	.00	.01	.00	.00	.35	.426	.820	1.220	.204	8.731
3.	.01	.00	.00	.00	.00	.00	.00	.00	.00	.00	.01	.00	.00	.01	.78	.480	.849	1.178	.166	9.671
4.	.20	.12	.00	.00	.00	.00	.00	.45	.00	.00	.00	.00	.00	.02	.02	.067	.894	1.118	.141	10.478
5.	.02	.02	.03	.00	.01	.05	.00	.00	.04	.00	.16	.23	.00	.15	.00	.116	.778	1.286	.110	11.867
6.	.00	.04	.00	.01	.00	.08	.00	.22	.05	.00	.00	.43	.01	.06	.00	.132	.586	1.705	.099	12.537
7.	.00	.00	.09	.01	.07	.01	.00	.00	.01	.00	.30	.05	.00	.30	.03	.624	.823	1.215	.083	13.651
8.	.05	.00	.00	.01	.00	.07	.23	.00	.05	.10	.04	.04	.00	.24	.00	.000	.902	1.108	.075	14.432
9.	.48	.63	.01	.00	.00	.00	.02	.00	.03	.02	.01	.03	.00	.00	.02	.682	.591	1.692	.072	14.661
10.	.06	.01	.05	.00	.04	.07	.14	.01	.01	.04	.27	.08	.12	.13	.01	.849	.820	1.219	.066	15.313
11.	.00	.00	.04	.01	.09	.02	.00	.01	.00	.00	.18	.00	.81	.03	.00	.491	.759	1.317	.050	17.550
12.	.00	.05	.61	.08	.24	.04	.07	.00	.05	.01	.02	.00	.00	.00	.00	.891	.880	1.137	.042	19.232
13.	.00	.01	.00	.28	.03	.00	.21	.00	.01	.61	.00	.01	.00	.00	.00	.034	.818	1.223	.036	20.705
14.	.02	.02	.05	.00	.11	.61	.00	.00	.71	.01	.00	.00	.00	.00	.01	.187	.908	1.101	.036	20.863
15.	.00	.00	.00	.39	.38	.04	.30	.00	.02	.16	.00	.02	.01	.00	.01	.298	.963	1.038	.032	21.865
16.	.02	.04	.11	.18	.01	.00	.02	.06	.00	.06	.01	.05	.02	.05	.04	.606	.933	1.072	.009	42.238

Note: Adjusted R Square = .029, Std. Error of the Estimate = .40190; Dependent Variable: ever lost any child(ren) aged 1-5 years; 1 = age of respondent, 2 = age of respondent at delivery, 3 = religion of respondent, 4 = marital status, 5 = husband had another wife, 6 = education of respondent, 7 = respondent occupation, 8 = number of living children 9 = spouse education, 10 = spouse occupation 11 = type of house living in, 12 = number of persons living in house, 13 = type of toilet facility for household, 14 = main source of water supply, 15 = distance of house to health facility, 16 = household waste disposal practices